



Continuity correction for barrier options in jump-diffusion models

El Hadj Aly Dia, Damien Lamberton

► To cite this version:

El Hadj Aly Dia, Damien Lamberton. Continuity correction for barrier options in jump-diffusion models. SIAM Journal on Financial Mathematics, 2011, 2 (1), pp.866-900. 10.1137/100817553 . hal-00547668

HAL Id: hal-00547668

<https://hal.science/hal-00547668>

Submitted on 17 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTINUITY CORRECTION FOR BARRIER OPTIONS IN JUMP-DIFFUSION MODELS

EL HADJ ALY DIA* AND DAMIEN LAMBERTON†

Abstract. The aim of this paper is to study the continuity correction for barrier options in jump-diffusion models. For this purpose, we express the pay-off of a barrier option in terms of the maximum of the underlying process. We then condition with respect to the jump times and to the values of the underlying at the jump times and rely on the connection between the maximum of the Brownian motion and Bessel processes.

Key words. Barrier option, Bessel process, Continuity correction, Exponential Lévy model, jump-diffusion.

AMS subject classifications. 60G51, 60J75, 91G20

JEL classification. C02, G13

1. Introduction. In the Black-Scholes setting, Broadie, Glasserman and Kou (1997) and Kou (2003) derived continuity correction formulas for barrier options. The purpose of this paper is to establish similar results for jump-diffusion models. The approach of Broadie, Glasserman and Kou was based on the expression of the pay-off of a barrier option in terms of the hitting time of the barrier by the underlying stock price. They managed to relate the discrete barrier option price to the continuous one by using classical results on the *overshoot* asymptotics of the Gaussian random walk.

Our approach is completely different and provides a new proof of the Broadie-Glasserman-Kou results, even in the Black-Scholes case. We start from the expression of the pay-off of barrier options in terms of the maximum process, which essentially involves the cumulative distribution function of the maximum. We then rely on the connection between the maximum of Brownian motion and the Bessel process, following the ideas of Asmussen, Glynn, Pitman (1995), in their study of the weak convergence of the normalized difference between the continuous and discrete maximum of Brownian motion. The extension to jump-diffusions is obtained by conditioning with respect to the jump times and to the values of the process at the jump times.

Note that the Asmussen-Glynn-Pitman Theorem was the basic tool for the derivation by Broadie, Glasserman and Kou (1999) of continuity corrections for *lookback* options, and we showed in [5] that this approach could be extended to jump-diffusion processes. The dependence of the payoff with respect to the maximum is much less smooth in the case of barriers, and we will need to go deeper into the connection between the maximum and the Bessel process to prove our results. In some sense, our results prove that continuity correction formulas can be obtained in a unified way for barrier and for lookback options.

The paper is organized as follows. In the next section, we present our main results: a continuity correction formula for a general pay-off (see Theorem 2.1), and its application to barrier options (see Proposition 2.2). We also demonstrate the use of these

*Université Paris-Est, Laboratoire d'Analyse et de Mathématiques Appliquées, UMR CNRS 8050, 5 bd. Descartes, Champs-sur-Marne, 77454 Marne-la-Vallée, France (dia.eha@gmail.com).

†Université Paris-Est, Laboratoire d'Analyse et de Mathématiques Appliquées, UMR CNRS 8050, 5 bd. Descartes, Champs-sur-Marne, 77454 Marne-la-Vallée, France (damien.lamberton@univ-mlv.fr).

results by showing some numerical results for a double-exponential jump-diffusion model. The other sections of the paper are devoted to the proof of Theorem 2.1. In Section 3, we derive some preliminary estimates on the jump times of a Poisson process. In Section 4, by conditioning with respect to the jump times, we reduce the problem to the analysis of discrete vs continuous supremum between the jump times.

In Section 5, we further condition with respect to the values of the underlying process at the jump times. We then have to deal with independent Brownian motions, and we establish a representation of a conditional expectation of a function of the maximum, the discrete maximum and the terminal value in terms of Bessel processes (see Proposition 5.2).

Section 6 is devoted to the derivation of some elementary estimates concerning the transition kernel of the Bessel process which are needed in the last two sections. In Section 7, we derive some bounds for conditional expectations, in order to be able to derive convergence results for the unconditional expectations from the corresponding results for conditional expectations. In Section 8, we establish the continuity correction for conditional expectations.

2. Continuity correction formulas. In a jump-diffusion model, the price of the underlying stock at time t is given by

$$S_t = S_0 e^{X_t}, \quad 0 \leq t \leq T,$$

where, under the pricing measure, the process $X = (X_t)_{0 \leq t \leq T}$ is given by

$$X_t = \gamma t + \sigma B_t + \sum_{i=1}^{N_t} Y_i, \quad (2.1)$$

where γ and σ are real constants, with $\sigma > 0$, $(B_t)_{0 \leq t \leq T}$ is a standard Brownian motion, N is a Poisson process with intensity $\lambda > 0$, and $(Y_i)_{i \geq 1}$ are i.i.d. random variables. Note that, under the pricing measure, the process $(e^{-(r-\delta)t} S_t)_{0 \leq t \leq T}$, where r is the interest rate and δ the dividend rate is a martingale. This implies the following relation between γ and the other parameters

$$\gamma = r - \delta - \frac{\sigma^2}{2} + \lambda \mathbb{E}(e^{Y_1} - 1).$$

In the terminology of exponential Lévy models, note that X is a Lévy process with a non-zero Brownian part and a finite Lévy measure, given by $\nu(dx) = \lambda \mu(dx)$, where μ is the distribution of the random variable Y_1 . For more details about Lévy processes see [12].

We define

$$M_t^X = \sup_{0 \leq s \leq t} X_s, \quad M_t^{X,n} = \max_{0 \leq k \leq n} X_{\frac{kt}{n}},$$

$$m_t^X = \inf_{0 \leq s \leq T} X_s, \quad m_t^{X,n} = \min_{0 \leq k \leq n} X_{\frac{kt}{n}}.$$

When there is no ambiguity we can remove the super index X .

The options we will consider in the sequel will have as underlying the asset with price S . We will denote by K and H the strike and the barrier of the option. The maturity of the options is assumed to be T . Figures 2.1 and 2.2 give the payoffs of barrier options. The corresponding prices are the expected values of the discounted payoffs.

Barrier	Continuous	Discrete
<i>Up Out</i>	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{M_T} < H\}}$	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{M_T^n} < H\}}$
<i>Up In</i>	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{M_T} \geq H\}}$	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{M_T^n} \geq H\}}$
<i>Down Out</i>	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{m_T} > H\}}$	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{m_T^n} > H\}}$
<i>Down In</i>	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{m_T} \leq H\}}$	$(S_T - K)^+ \mathbb{1}_{\{S_0 e^{m_T^n} \leq H\}}$

FIG. 2.1. Payoffs of barrier call options

Barrier	Continuous	Discrete
<i>Up Out</i>	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{M_T} < H\}}$	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{M_T^n} < H\}}$
<i>Up In</i>	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{M_T} \geq H\}}$	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{M_T^n} \geq H\}}$
<i>Down Out</i>	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{m_T} > H\}}$	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{m_T^n} > H\}}$
<i>Down In</i>	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{m_T} \leq H\}}$	$(K - S_T)^+ \mathbb{1}_{\{S_0 e^{m_T^n} \leq H\}}$

FIG. 2.2. Payoffs of barrier put options

Let $UOC(H)$ be the price of a continuous *up and out* call with barrier H , We have

$$UOC(H) = \mathbb{E} e^{-rT} (S_0 e^{X_T} - K)^+ \mathbb{1}_{\{\sup_{0 \leq t \leq T} S_0 e^{X_t} < H\}}.$$

Define

$$k = \log \left(\frac{K}{S_0} \right), \quad h = \log \left(\frac{H}{S_0} \right).$$

We can write

$$\begin{aligned} UOC(H) &= S_0 \mathbb{E} e^{-rT} e^{X_T} \mathbb{1}_{\{M_T < h, X_T > k\}} - K e^{-rT} \mathbb{P}[M_T < h, X_T > k] \\ &= S_0 e^{-\delta T} \mathbb{E} e^{-(r-\delta)T} e^{X_T} \mathbb{1}_{\{M_T < h, X_T > k\}} - K e^{-rT} \mathbb{P}[M_T < h, X_T > k]. \end{aligned}$$

We know that the process $(e^{-(r-\delta)t} e^{X_t})_{0 \leq t \leq T}$ is a martingale. Let $\bar{\mathbb{P}}$ be the probability defined by its density with respect to the pricing probability measure \mathbb{P}

$$\frac{d\bar{\mathbb{P}}}{d\mathbb{P}} = e^{-(r-\delta)T + X_T}.$$

Note that (as can be deduced, for instance, from Theorem 3.9 of [10]), the process X remains a Lévy process under probability $\bar{\mathbb{P}}$, and that its Lévy measure under $\bar{\mathbb{P}}$ is given by $\bar{\nu}(dx) = e^x \nu(dx)$.

We have

$$UOC(H) = S_0 e^{-\delta T} \bar{\mathbb{P}}[M_T < h, X_T > k] - K e^{-rT} \mathbb{P}[M_T < h, X_T > k]$$

If we call UOC^n the price of a discrete *up and out* call with barrier H , and n fixing dates (with step $\frac{T}{n}$), then we have similarly

$$UOC^n(H) = S_0 e^{-\delta T} \bar{\mathbb{P}}[M_T^n < h, X_T > k] - K e^{-rT} \mathbb{P}[M_T^n < h, X_T > k].$$

Finding continuity corrections between continuous and discrete barrier options amounts in fact to finding corrections between the above probabilities. This is the aim of the following result.

THEOREM 2.1. *Let X be an integrable Lévy process of the form (2.1), with $\sigma > 0$. For any bounded Borel measurable function $g : \mathbb{R} \rightarrow \mathbb{R}$ and for any positive number x , we have*

$$\mathbb{E} \left[g(X_T) \mathbf{1}_{\{M_T \geq x > M_T^n\}} \right] = \mathbb{E} \left[g(X_T) \mathbf{1}_{\{M_T \geq x > M_T - \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}\}} \right] + o\left(\frac{1}{\sqrt{n}}\right),$$

where $\beta_1 = \mathbb{E}R$ and R is defined by

$$R = \min_{\{j \in \mathbb{Z}\}} \check{R}(U + j). \quad (2.2)$$

Here, $(\check{R}(t))_{t \in \mathbb{R}}$ is a two sided three dimensional Bessel process (i.e. $\check{R}(t) = R_1(t)$ for $t \geq 0$ and $\check{R}(t) = R_2(-t)$ for $t < 0$, where R_1 and R_2 are independent copies of the usual three dimensional Bessel process, starting from 0) and U is uniformly distributed on $[0, 1]$ and independent of \check{R} .

Note that the result does not depend on the jump part of the process, so that the continuity correction for jump-diffusion models is the same as for the Black-Scholes model.

The result of Theorem 2.1 can also be written in the form

$$\mathbb{E} \left[g(X_T) \mathbf{1}_{\{M_T < x + \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}\}} \right] = \mathbb{E} \left[g(X_T) \mathbf{1}_{\{M_T^n < x\}} \right] + o\left(\frac{1}{\sqrt{n}}\right).$$

Moreover, the proof of Theorem 2.1 shows that the theorem is still true if we replace x by a sequence x_n which converges to x when $n \rightarrow +\infty$. So, under the assumptions of Theorem 2.1, we have

$$\mathbb{P} \left(M_T < x + \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}, X_T > y \right) = \mathbb{P} (M_T^n < x, X_T > y) + o\left(\frac{1}{\sqrt{n}}\right) \quad (2.3)$$

$$\mathbb{P} (M_T < x, X_T > y) = \mathbb{P} \left[M_T^n < x - \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}, X_T > y \right] + o\left(\frac{1}{\sqrt{n}}\right) \quad (2.4)$$

Therefore, we deduce from Theorem 2.1 the relations between continuous and discrete barrier options.

PROPOSITION 2.2. *Let X be a Lévy process with generating triplet (γ, σ^2, ν) satisfying $\sigma > 0$ and $\nu(\mathbb{R}) < \infty$, $V(H)$ be the price of a continuous option with barrier H , and $V^n(H)$ be the price of the corresponding discrete barrier option. We assume that the process $(e^{X_t - (r-\delta)t})_{t \geq 0}$ is a martingale. Then*

$$\begin{aligned} V^n(H) &= V \left(H e^{\pm \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}} \right) + o\left(\frac{1}{\sqrt{n}}\right) \\ V(H) &= V^n \left(H e^{\mp \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}} \right) + o\left(\frac{1}{\sqrt{n}}\right), \end{aligned}$$

where in \pm and \mp , the top case applies for Up options and the bottom case applies for Down options.

REMARK 2.3. Under the assumptions of Proposition 2.2, we can prove that

$$V(H) - V^n(H) = \frac{C}{\sqrt{n}} + o\left(\frac{1}{\sqrt{n}}\right).$$

See Remark 8.7.

Proof of Proposition 2.2. For the proof, we will consider only barrier options without *rebate*, since the latter can be easily deduced from the former. Theorem 2.1 and (2.3) lead obviously to the following results

$$\begin{aligned} \mathbb{P}\left[M_T < y + \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}, X_T \leq x\right] &= \mathbb{P}[M_T^n < y, X_T \leq x] + o\left(\frac{1}{\sqrt{n}}\right) \\ \mathbb{P}\left[M_T \geq y + \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}, X_T \leq x\right] &= \mathbb{P}[M_T^n \geq y, X_T \leq x] + o\left(\frac{1}{\sqrt{n}}\right) \\ \mathbb{P}\left[M_T < y + \frac{\sigma\sqrt{T}\beta_1}{\sqrt{n}}, X_T < x\right] &= \mathbb{P}[M_T^n < y, X_T < x] + o\left(\frac{1}{\sqrt{n}}\right). \end{aligned}$$

The price of barrier options can be written in terms of the above probabilities (as in the case of the call *Up and Out* studied in the beginning of the section). Recall that in the *Down* case, the infimum process m of X satisfies

$$\begin{aligned} m_t &= \inf_{0 \leq s \leq t} X_s \\ &= - \sup_{0 \leq s \leq t} (-X_s). \end{aligned}$$

We deduce the first result of the proposition. For the second part of the proposition, we proceed in the same way and use (2.4). \diamond

We will test the performance of Proposition 2.2 with the double exponential *jump-diffusion* model (see [8]). So, we have

$$X_s = \gamma s + \sigma B_s + \sum_{i=1}^{N_s} Y_i,$$

where N is a poisson process with intensity λ , and Y_1 follows an asymmetric double exponential distribution with probability density function

$$f_Y(y) = p\eta_1 e^{-\eta_1 y} \mathbb{1}_{\{y \geq 0\}} + q\eta_2 e^{\eta_2 y} \mathbb{1}_{\{y < 0\}},$$

where η_1, η_2 are positive numbers (with $\eta_1 > 1$ to ensure integrability of the exponential), and the non-negative real numbers p and q satisfy $p + q = 1$. In our numerical examples, the values of the parameters are the following: $\sigma = 0.3$, $p = 0.6$, $\lambda = 7$, $\eta_1 = 50$ and $\eta_2 = 25$. We will consider the *up and out put* option with parameters $S_0 = 100$, $r = 0.05$, $\delta = 0$, $T = 1$, $K = 100$, $H = 110$ and *rebate* = 10. The continuous price, computed by the method used in [9], is equal to 13.240. The discrete prices are computed by Monte Carlo methods. In Table 2.1, we study the convergence of the discrete price and the corrected discrete price (using the second equality in Proposition 2.2) to the continuous price.

n	Discrete price	Relative error	Corrected discrete price	Relative error
5	14.193	7.201%	13.883	4.857%
6	14.160	6.945%	13.772	4.016%
7	14.128	6.707%	13.667	3.379%
8	14.095	6.459%	13.627	2.923%
9	14.072	6.278%	13.577	2.544%
10	14.048	6.101%	13.542	2.273%
15	13.957	5.410%	13.429	1.430%
25	13.851	4.619%	13.358	0.896%

TABLE 2.1

Performance of the continuity correction in double exponential jump-diffusion model.

As expected, the discrete price converges slowly, while the corrected price converges rapidly to the continuous price. The reverse problem is studied in Table 2.2. We approximate the discrete barrier price by the corrected continuous price according to our correction formula (see the first result of Proposition 2.2). In the last column we give the relative error made by approximating the discrete price by the corrected continuous price. The latter clearly is a good approximation of the discrete price, compared to the continuous price.

n	Discrete price	Corrected continuous price	Relative Error
5	14.193	13.964	1.613%
10	14.048	13.894	1.096%
15	13.957	13.829	0.917%
20	13.896	13.780	0.834%
25	13.851	13.742	0.787%
30	13.816	13.711	0.760%
35	13.789	13.685	0.754%
40	13.764	13.664	0.726%
45	13.743	13.645	0.713%
50	13.726	13.629	0.707%

TABLE 2.2

Performance of the continuity correction in double exponential jump-diffusion model.

3. Estimates for the Poisson process. In this section, we give some estimates for the jump times of a Poisson process. These estimates will be used to derive domination conditions in order to justify the convergence of some expectations.

PROPOSITION 3.1. *Let $(N_t)_{t \geq 0}$ be a homogeneous Poisson process, with jump times $(T_l)_{l \geq 1}$. For $t > 0$ fixed and for any integer $l \geq 1$, we have, for $i = 1, \dots, l$,*

$$\mathbb{E} \left(\frac{1}{\sqrt{T_i - T_{i-1}}} \mid N_t = l \right) \leq \frac{2l}{\sqrt{t}}$$

and

$$\mathbb{E} \left(\frac{1}{\sqrt{t - T_l}} \mid N_t = l \right) \leq \frac{2l}{\sqrt{t}}.$$

Proof. Using the conditional distribution of the jump times T_1, \dots, T_l , given $\{N_t = l\}$, we have

$$\begin{aligned} \mathbb{E} \left(\frac{1}{\sqrt{T_i - T_{i-1}}} \mid N_t = l \right) &= \int_{\{0 < t_1 < \dots < t_l < t\}} \frac{1}{\sqrt{t_i - t_{i-1}}} \frac{l!}{t^l} dt_1 \dots dt_l \\ &= \int_{\{u_1 > 0, \dots, u_l > 0, \sum_{j=1}^l u_j < t\}} \frac{1}{\sqrt{u_i}} \frac{l!}{t^l} du_1 \dots du_l \\ &\leq \frac{2l}{\sqrt{t}} \int_{\{u_1 > 0, \dots, u_l > 0, \sum_{j \neq i} u_j < t\}} \frac{(l-1)!}{t^{l-1}} du_1 \dots \cancel{du_i} \dots du_l \\ &= \frac{2l}{\sqrt{t}}, \end{aligned}$$

where we have used $\int_0^t \frac{du_i}{\sqrt{u_i}} = 2/\sqrt{t}$ and, in the last integral, u_i is omitted. The proof of the second inequality is similar. \diamond

PROPOSITION 3.2. *Let $(N_t)_{t \geq 0}$ be a homogeneous Poisson process, with jump times $(T_l)_{l \geq 1}$. For $t > 0$ fixed and for any integer $l \geq 1$, we have, for $i = 1, \dots, l$ and for any $\alpha > 0$,*

$$\mathbb{P}(T_i - T_{i-1} \leq \alpha t \mid N_t = l) \leq l\alpha$$

and

$$\mathbb{P}(t - T_l \leq \alpha t \mid N_t = l) \leq l\alpha.$$

Proof. We can assume that $\alpha < 1$ and write, for $i = 1, \dots, l$, using the conditional distribution of jump times given $\{N_t = l\}$,

$$\begin{aligned} \mathbb{P}(T_i - T_{i-1} \leq \alpha t \mid N_t = l) &= \int_{\{0 < t_1 < \dots < t_l < t\}} \mathbb{1}_{\{t_i - t_{i-1} \leq \alpha t\}} \frac{l!}{t^l} dt_1 \dots dt_l \\ &= \int_{\{u_1 > 0, \dots, u_l > 0, \sum_{j=1}^l u_j < t\}} \mathbb{1}_{\{u_i \leq \alpha t\}} \frac{l!}{t^l} du_1 \dots du_l \\ &\leq l\alpha \int_{\{u_1 > 0, \dots, u_l > 0, \sum_{j \neq i} u_j < t\}} \frac{(l-1)!}{t^{l-1}} du_1 \dots \cancel{du_i} \dots du_l \\ &= l\alpha, \end{aligned}$$

where, in the last integral, the variable u_i is omitted. The proof of the second inequality is similar. \diamond

4. Conditioning with respect to the jump times. For the proof of Theorem 2.1, we will first condition with respect to the jump times of the Poisson process. Fix $x > 0$ and $t > 0$. We have

$$\begin{aligned} \mathbb{E}(g(X_t) \mathbb{1}_{\{M_t \geq x\}}) - \mathbb{E}(g(X_t) \mathbb{1}_{\{M_t^n \geq x\}}) &= \mathbb{E}(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}}) \\ &= \sum_{l=0}^{\infty} \mathbb{E}(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \mid N_t = l) \mathbb{P}(N_t = l). \end{aligned}$$

Conditionally on $\{N_t = 0\}$, we have $X_s = \gamma s + \sigma B_s$, for $s \in [0, t]$, and

$$\mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \mid N_t = 0 \right) = \mathbb{E} \left(g(X_t) \mathbb{1}_{\{M^0 \geq x > M^{0,n}\}} \mid N_t = 0 \right),$$

with $M^0 = \sup_{0 \leq s \leq t} (\gamma s + \sigma B_s)$, $M^{0,n} = \max_{k=0, \dots, n} X_{kt/n}$.

Conditionally on $\{N_t = l\}$ and $\{T_1 = t_1, \dots, T_l = t_l\}$, with $0 < t_1 < \dots < t_l < t$, we set $t_{l+1} = t$ and, for $j = 0, \dots, l$,

$$M^j = \sup_{s \in [t_j, t_{j+1})} X_s, \quad M^{j,n} = \max_{k, kt/n \in [t_j, t_{j+1})} X_{kt/n},$$

with, by convention $M^{j,n} = -\infty$ if there is no integer k such that $kt/n \in [t_j, t_{j+1})$. In the sequel, we denote by θ the vector (t_1, \dots, t_l) and by $\mathbb{E}_{l,\theta}$ the conditional expectation given $\{N_t = l, T_1 = t_1, \dots, T_l = t_l\}$. Conditionally on the values of X at times t_j , the random variables M^j are independent and have a density. So they are almost surely pairwise distinct and we have

$$\begin{aligned} \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) &= \sum_{j=0}^l \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n, M^j > \max_{i \neq j} M^i\}} \right) \\ &= \sum_{j=0}^l \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M_t^n, M^j > \max_{i \neq j} M^i\}} \right). \end{aligned}$$

Hence

$$\mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) = \sum_{j=0}^l \left(\alpha_l^{j,n}(\theta) - \beta_l^{j,n}(\theta) \right),$$

with

$$\alpha_l^{j,n}(\theta) = \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}} \right)$$

and

$$\beta_l^{j,n}(\theta) = \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i, \max_{i \neq j} M^{i,n} \geq x\}} \right).$$

Integrating with respect to the jump times, we get

$$\mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) = \mathbb{E} \left(\alpha_{N_t}^n(T_1, \dots, T_{N_t}) - \beta_{N_t}^n(T_1, \dots, T_{N_t}) \right),$$

where for $l \in \mathbb{N}$,

$$\alpha_l^n(t_1, \dots, t_l) = \mathbb{1}_{\{l \geq 1\}} \sum_{j=0}^l \alpha_l^{j,n}(t_1, \dots, t_l)$$

and

$$\beta_l^n(t_1, \dots, t_l) = \mathbb{1}_{\{l \geq 1\}} \sum_{j=0}^l \beta_l^{j,n}(t_1, \dots, t_l).$$

With these notations, we can state the following proposition.

PROPOSITION 4.1. *We have $\lim_{n \rightarrow +\infty} \sqrt{n} \mathbb{E} \left(\beta_{N_t}^n(T_1, \dots, T_{N_t}) \right) = 0$.*

For the proof of this proposition, we will use the following reformulation of the Asmussen-Glynn-Pitman Theorem. It can be deduced from a careful reading of the proof of Theorem 1 in [1] (see particularly pages 879 to 883, and Remark 2).

THEOREM 4.2. *Consider four real numbers a, b, x and y , with $0 \leq a < b$. Let $\beta = (\beta_s)_{a \leq s \leq b}$ be a Brownian bridge from x to y over the time interval $[a, b]$ (so that $\beta_a = x$ and $\beta_b = y$) and let t be a fixed positive number. Denote by M the supremum of β and, for any positive integer n , by M^n the discrete supremum associated with a mesh of size t/n , so that*

$$M = \sup_{a \leq s \leq b} \beta_s \quad \text{and} \quad M^n = \sup_{k \in I_n} \beta_{\frac{kt}{n}}, \quad \text{where } I_n = \left\{ k \in \mathbb{N} \mid \frac{kt}{n} \in [a, b] \right\}.$$

Then, as n goes to infinity, the pair $(\sqrt{n}(M - M^n), \beta)$ converges in distribution to the pair $(\sqrt{t}R, \beta)$ where R , defined in Theorem 2.1, is independent of β .

Proof of Proposition 4.1. We have

$$\begin{aligned} |\beta_l^{j,n}(\theta)| &\leq \|g\|_\infty \sum_{i \neq j} \mathbb{P}_{l,\theta} (M^j \geq x > M^{j,n}, M^j > M^i \geq M^{i,n} \geq x) \\ &\leq \|g\|_\infty \sum_{i \neq j} \mathbb{P}_{l,\theta} (x \leq M^j < x + (M^j - M^{j,n}), x \leq M^i < x + (M^j - M^{j,n})). \end{aligned}$$

Conditionally on $\{N_t = l\}$, $\{(T_1, \dots, T_l) = \theta\}$ and $\{X_{T_k} = x_k, k = 1, \dots, l\}$, the processes $(X_s - X_{t_i})_{t_i \leq s < t_{i+1}}$ and $(X_s - X_{t_j})_{t_j \leq s < t_{j+1}}$ (for $i \neq j$) are independent Brownian motions. The pairs of random variables $(M^j - x_j, M^j - M^{j,n})$ and $(M^i - x_i, M^i - M^{i,n})$ are independent and we have

$$\mathbb{E}(\mathbf{1}_{\{x \leq M^i < x + M^j - M^{j,n}\}} \mid M^j, M^{j,n}) = \int_{x-x_i}^{x-x_i+M^j-M^{j,n}} f_i(u) du,$$

where f_i is the probability density function of the random variable $\sup_{0 \leq s \leq t_{i+1}-t_i} (\gamma s + \sigma B_s)$. We know (see for example Lemma 2.22 of [4]) that the function f_i is bounded by $C/\sqrt{t_{i+1}-t_i}$, where the constant C depends only on γ, σ and t . We deduce that

$$\begin{aligned} &\mathbb{P}_{l,\theta} (x \leq M^j < x + (M^j - M^{j,n}), x \leq M^i < x + (M^j - M^{j,n})) \\ &\leq \frac{C}{\sqrt{t_{i+1}-t_i}} \mathbb{E}_{l,\theta} ((M^j - M^{j,n}) \mathbf{1}_{\{x \leq M^j < x + (M^j - M^{j,n})\}}), \end{aligned}$$

Note that by Theorem 4.2 and the fact that M^j has a continuous distribution, the sequence $(\sqrt{n}(M^j - M^{j,n}) \mathbf{1}_{\{x \leq M^j < x + (M^j - M^{j,n})\}})_{n \in \mathbb{N}}$ converges to 0 in probability and, since $(\sqrt{n}(M^j - M^{j,n}))_{n \in \mathbb{N}}$ is uniformly integrable (see [1], Lemma 6), we have

$$\lim_{n \rightarrow +\infty} \sqrt{n} \mathbb{E}_{l,\theta} ((M^j - M^{j,n}) \mathbf{1}_{\{x \leq M^j < x + (M^j - M^{j,n})\}}) = 0.$$

On the other hand, we have

$$\sum_{j=0}^l \beta_l^{j,n}(\theta) \leq \sum_{j=0}^l \sum_{i \neq j} \frac{C}{\sqrt{t_{i+1}-t_i}} \sqrt{\frac{t}{n}} \quad (4.1)$$

$$\leq Cl \sqrt{\frac{t}{n}} \sum_{i=0}^l \frac{1}{\sqrt{t_{i+1}-t_i}}. \quad (4.2)$$

We deduce that the sequence of random variables $\sqrt{n}\beta_{N_t}^n(T_1, \dots, T_{N_t})$ is dominated by an integrable random variable by Proposition 3.1. This concludes the proof. \diamond

5. Conditioning with respect to the positions at jump times and representation using the Bessel process. It follows from the discussion of Section 4 that

$$\begin{aligned} \mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) &= \mathbb{E}_0 \left(g(X_t) \mathbb{1}_{\{M^0 \geq x > M_t^{0,n}\}} \right) \mathbb{P}(N_t = 0) \\ &\quad + \mathbb{E} \left(\alpha_{N_t}^n(T_1, \dots, T_{N_t}) \right) + o(1/\sqrt{n}), \end{aligned}$$

where $\mathbb{E}_0 = \mathbb{E}(\cdot \mid N_t = 0)$ and $\alpha_l^n(t_1, \dots, t_l) = \sum_{j=0}^l \alpha_l^{j,n}(t_1, \dots, t_l)$. We have

$$\alpha_l^{j,n}(t_1, \dots, t_l) = \mathbb{E} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}} \mid N_t = l, T_1 = t_1, \dots, T_l = t_l \right).$$

For $j = 0, \dots, l$, we set

$$\hat{\beta}_u^j = \frac{X_{t_j+u(t_{j+1}-t_j)}}{\sigma\sqrt{t_{j+1}-t_j}}, \text{ for } u \in [0, 1), \quad \text{and } \hat{\beta}_1^j = \frac{X_{t_{j+1}}^-}{\sigma\sqrt{t_{j+1}-t_j}}.$$

We have, putting $\sigma_j = \sigma\sqrt{t_{j+1}-t_j}$,

$$M^j = \sigma_j \hat{M}^j \quad \text{and} \quad M^{j,n} = \sigma_j \hat{M}^{j,n},$$

where

$$\hat{M}^j = \sup_{u \in [0,1]} \hat{\beta}_u^j \quad \text{and} \quad \hat{M}^{j,n} = \sup_{k \in I_n^j} \hat{\beta}_{\frac{\lambda_{j,k}}{n} - t_j}^j,$$

with the notations $\lambda_j = t/(t_{j+1}-t_j)$, $\hat{t}_j = t_j/(t_{j+1}-t_j)$ and

$$I_n^j = \{k \in \mathbb{N} \mid t_j \leq kt/n < t_{j+1}\} \text{ for } j = 0, \dots, l-1$$

and $I_n^l = \{k \in \mathbb{N} \mid t_j \leq kt/n \leq t_{l+1} = t\}$. Here again we use the convention $\hat{M}^{j,n} = -\infty$ if $I_n^j = \emptyset$.

For the computation of $\alpha_l^{j,n}(t_1, \dots, t_l)$, we will further condition with respect to $\{X_{T_1} = x_1, \dots, X_{T_l} = x_l\}$, where x_1, \dots, x_l are arbitrary real numbers. So, we introduce the notations

$$\theta = (t_1, \dots, t_l), \quad \xi = (x_1, \dots, x_l),$$

and

$$\mathbb{P}_{l,\theta,\xi} = \mathbb{P}(\cdot \mid N_t = l, T_k = t_k, X_{T_k} = x_k, k = 1, \dots, l).$$

The expectation under $\mathbb{P}_{l,\theta,\xi}$ will be denoted by $\mathbb{E}_{l,\theta,\xi}$. Note that, under $\mathbb{P}_{l,\theta,\xi}$, the processes $\hat{\beta}^j$ are independent Brownian motions on the interval $[0, 1]$, with initial values $\hat{\beta}_0^j = \hat{x}_j$, with $\hat{x}_j = x_j/\sigma_j$, $j = 0, \dots, l$.

With these notations, we can state the following lemma.

LEMMA 5.1. *We have*

$$\mathbb{E}_{l,\theta,\xi} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}} \right) = \mathbb{E}_{l,\theta,\xi} \left(\mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \alpha_{l,\theta,\xi}^j(X_{t_{j+1}}^-, M^j) \right),$$

where

$$\alpha_{l,\theta,\xi}^j(y, m) = \begin{cases} \mathbb{E}_{l,\theta,\xi}(g(X_t) \mathbb{1}_{\{\max_{i \neq j} M^i < m\}}) & \text{if } j = 0, \dots, l-1 \\ g(y) \mathbb{P}_{l,\theta,\xi}(\max_{i \neq j} M^i < m) & \text{if } j = l. \end{cases}$$

Proof. Note that, if $j < l$, under $\mathbb{P}_{l,\theta,\xi}$, the pair $(M^j, M^{j,n})$ on the one hand, and the random variables X_t, M^i for $i \neq j$ on the other hand are independent, so that

$$\mathbb{E}_{l,\theta,\xi}(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}}) = \mathbb{E}_{l,\theta,\xi}(\mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \alpha_{l,\theta,\xi}^j(M^j)),$$

where

$$\alpha_{l,\theta,\xi}^j(m) = \mathbb{E}_{l,\theta,\xi}(g(X_t) \mathbb{1}_{\{\max_{i \neq j} M^i < m\}}).$$

Note that in this case the function $\alpha_{l,\theta,\xi}^j$ does not depend on y . Now, if $j = l$, we have $X_t = X_{t_{j+1}}^-$ and the random variables M^i for $i < l$ are independent of the pair $(X_{t_{l+1}}, M^l)$, so that

$$\mathbb{E}_{l,\theta,\xi}(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}}) = \mathbb{E}_{l,\theta,\xi}(\mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \alpha_{l,\theta,\xi}^j(X_{t_{j+1}}, M^j)),$$

with

$$\alpha_{l,\theta,\xi}^j(y, m) = g(y) \mathbb{P}_{l,\theta,\xi}\left(\max_{i \neq j} M^i < m\right).$$

◇

We will now give a representation of the expectations in Lemma 5.1 in terms of Bessel processes. Set $\tau^j = \sup\{u \in [0, 1] \mid \hat{\beta}_u^j = \hat{M}^j\}$. Conditionally on $\tau^j = s$ and $\hat{M}^j = m$, we set $R_1^j(u) = m - \hat{\beta}_{s-u}^j$, for $u \in [0, s]$ and $R_2^j(v) = m - \hat{\beta}_{s+v}^j$, for $v \in [0, 1-s]$. We know that, conditionally on $\{\tau^j = s, \hat{M}^j = m, \hat{\beta}_1^j = y\}$, the processes R_1^j et R_2^j are independent Bessel bridges of dimension 3 (cf. [1], Proposition 2). We can write, conditionally on $\{\tau^j = s, \hat{M}^j = m\}$,

$$\hat{M}^j - \hat{M}^{j,n} = \min_{k \in I_n^-} R_1^j(s + \hat{t}_j - \lambda_j(k/n)) \wedge \min_{k \in I_n^+} R_2^j(\lambda_j(k/n) - \hat{t}_j - s),$$

with

$$I_n^- = \{k \in \mathbb{N} \mid 0 \leq s + \hat{t}_j - \lambda_j(k/n) \leq s\}, \quad I_n^+ = \{k \in \mathbb{N} \mid s \leq \lambda_j(k/n) - \hat{t}_j \leq 1\}.$$

Hence

$$\hat{M}^j - \hat{M}^{j,n} = \min_{0 \leq k \leq N_n^1} R_1^j(d_n^j(s) + \lambda_j(k/n)) \wedge \min_{1 \leq k \leq N_n^2} R_2^j(\lambda_j(k/n) - d_n^j(s)), \quad (5.1)$$

with $d_n^j(s) = \hat{t}_j + s - \frac{\lambda_j}{n} \left\lceil \frac{n(\hat{t}_j + s)}{\lambda_j} \right\rceil$ (where $[x]$ is the greatest integer in x ; note that $0 \leq d_n^j(s) \leq \lambda_j/n$) and

$$\begin{aligned} N_n^1 &= \max\{k \in \mathbb{N} \mid d_n^j(s) + \lambda_j(k/n) \leq s\}, \\ N_n^2 &= \max\{k \in \mathbb{N} \mid -d_n^j(s) + \lambda_j(k/n) \leq 1 - s\}. \end{aligned}$$

Note that N_n^1 is a well defined non-negative integer if I_n^- is not empty and N_n^2 is a well defined positive integer if I_n^+ is not empty. When one of the two sets is empty and not the other, the minimum in (5.1) is considered on the non-empty set. Note that, if $\lambda_j/n < 1$ (or, equivalently, $t_{j+1} - t_j > t/n$), at least one of the two sets is non-empty. In the following proposition, we will also use the notation

$$\gamma_j = \gamma \sqrt{t_{j+1} - t_j} / \sigma.$$

It should be emphasized that, in the next statement, there is no conditioning on the terminal values of the Bessel processes, in contrast with the statement of Proposition 2 of [1].

PROPOSITION 5.2. *Assume $\lambda_j/n < 1$. For any bounded Borel measurable function $F : \mathbb{R}^3 \rightarrow \mathbb{R}$, we have*

$$\mathbb{E}_{l,\theta,\xi} F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) = \int_0^1 ds \mathbb{E} \left(\hat{F} \left(R_1(s), R_2(1-s), R_s^{j,n} \right) \right),$$

where

$$\hat{F}(r_1, r_2, \rho) = \frac{1}{2} \frac{e^{\gamma_j(r_1-r_2) - (\gamma_j^2/2)}}{r_1 r_2} F(x_j + \sigma_j(r_1 - r_2), x_j + \sigma_j r_1, \sigma_j \rho),$$

R_1 et R_2 are independent three-dimensional Bessel processes, starting from 0, and

$$R_s^{j,n} = \min_{-N_n^2 \leq k \leq N_n^1} \check{R}(d_n^j(s) + \lambda_j(k/n)),$$

with $\check{R}(u) = R_1(u)$ for $u \geq 0$ and $\check{R}(u) = R_2(-u)$ for $u < 0$.

Proof. Note that $F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) = F \left(\sigma_j \hat{\beta}_1, \sigma_j \hat{M}^j, \sigma_j (\hat{M}^j - \hat{M}^{j,n}) \right)$. In view of the discussion before the statement of Proposition 5.2, we observe that the conditional distribution of $\hat{M}^j - \hat{M}^{j,n}$ given $\{\tau^j = s, \hat{M}^j = m, \hat{\beta}_1^j = y\}$ is the same as the conditional distribution of $R_s^{j,n}$ given $\{R_1(s) = m - \hat{x}_j, R_2(1-s) = m - y\}$, so that

$$\begin{aligned} \mathbb{E}_{l,\theta,\xi} \left(F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) \mid \tau^j = s, \hat{M}^j = m, \hat{\beta}_1^j = y \right) &= \\ \mathbb{E} \left(F \left(\sigma_j y, \sigma_j m, \sigma_j R_s^{j,n} \right) \mid R_1(s) = m - \hat{x}_j, R_2(1-s) = m - y \right) &= \\ \psi_s^j(m - \hat{x}_j, m - y), \end{aligned}$$

with

$$\psi_s^j(r_1, r_2) = \mathbb{E} \left(G(r_1, r_2, R_s^{j,n}) \mid R_1(s) = r_1, R_2(1-s) = r_2 \right),$$

and

$$G(r_1, r_2, \rho) = F(x_j + \sigma_j(r_1 - r_2), x_j + \sigma_j r_1, \sigma_j \rho).$$

Recall that, under probability $\mathbb{P}_{l,\theta,\xi}$, the process $(\hat{\beta}_u^j)_{u \in [0,1]}$ is a Brownian motion, starting from $\hat{x}_j = x_j/\sigma_j$, with drift γ_j and with unit variance coefficient. It follows that the conditional probability density function of the pair (τ^j, \hat{M}^j) given $\hat{\beta}_1^j = y$ can be written

$$\mathbb{P}(\tau^j \in ds, \hat{M}^j \in dm \mid \hat{\beta}_1^j = y) = \frac{u(s, m - \hat{x}_j)u(1-s, m - y)}{n(y - \hat{x}_j)} ds dm,$$

where n is the probability density function of the standard normal distribution and

$$u(s, m) = \frac{1}{\sqrt{\pi}} \frac{m}{s^{3/2}} e^{-m^2/(2s)}, \quad m > 0.$$

The above expression of the conditional distribution of (τ^j, \hat{M}^j) follows from Proposition 8.15 in Chapter II of [6].

We now have

$$\begin{aligned} \mathbb{E}_{l, \theta, \xi} F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) &= \\ &= \int \mathbb{P}(\tau^j \in ds, \hat{M}^j \in dm, \hat{\beta}_1^j \in dy) \psi_s^j(m - \hat{x}_j, m - y) \\ &= \int \mathbb{P}(\hat{\beta}_1^j \in dy) \int_0^1 ds \int_{\hat{x}_j \vee y}^\infty \frac{dm}{n(y - \hat{x}_j)} u(s, m - \hat{x}_j) u(1 - s, m - y) \psi_s^j(m - \hat{x}_j, m - y). \end{aligned}$$

Since $\mathbb{P}(\hat{\beta}_1^j \in dy) = n(y - \hat{x}_j - \gamma_j) dy$, we can write, with the substitution $r_1 = m - \hat{x}_j$, $r_2 = m - y$ in the integral with respect to y and m ,

$$\begin{aligned} \mathbb{E}_{l, \theta, \xi} F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) &= \\ &= \int_0^1 ds \int_0^\infty dr_1 \int_0^\infty dr_2 e^{\gamma_j(r_1 - r_2) - (\gamma_j^2/2)} u(s, r_1) u(1 - s, r_2) \psi_s^j(r_1, r_2). \end{aligned} \quad (5.2)$$

Recall that the transition density of the three-dimensional Bessel process is given by

$$\tilde{q}_t(x, y) = \frac{1}{x} q_t(x, y) y, \quad x, y > 0, t > 0,$$

where $q_t(x, y)$ is the transition density of Brownian motion (on $[0, +\infty)$) killed when it hits 0, which can be written

$$q_t(x, y) = g_t(x - y) - g_t(x + y),$$

where g_t is the density of the normal distribution with mean 0 and variance t . For these properties of the Bessel process, see [11] (Chapter VI, Section 3). For $x = 0$, we have

$$\tilde{q}_t(0, y) = \sqrt{\frac{2}{\pi}} \frac{y^2}{t^{3/2}} e^{-y^2/(2t)}, \quad y > 0, \quad t > 0.$$

Note that, for any $m > 0$ and for any $s > 0$,

$$u(s, m) = \frac{1}{m\sqrt{2}} \tilde{q}_s(0, m).$$

Hence, (5.2) can be written as follows

$$\begin{aligned} \mathbb{E}_{l, \theta, \xi} F \left(X_{t_{j+1}^-}, M^j, M^j - M^{j,n} \right) &= \\ &= \frac{1}{2} \int_0^1 ds \int_0^\infty \int_0^\infty \mathbb{P}(R_1^j(s) \in dr_1, R_2^j(1 - s) \in dr_2) \frac{e^{\gamma_j(r_1 - r_2) - (\gamma_j^2/2)}}{r_1 r_2} \psi_s^j(r_1, r_2) \\ &= \int_0^1 ds \mathbb{E} \left(L_s^j \mathbb{E} \left(G(R_1(s), R_2(1 - s), R_s^{j,n}) \mid R_1(s), R_2(1 - s) \right) \right), \end{aligned}$$

where

$$L_s^j = \frac{e^{\gamma_j(R_1(s) - R_2(1-s)) - (\gamma_j^2/2)}}{2R_1(s)R_2(1-s)}. \quad (5.3)$$

The proposition then follows from the equality

$$\hat{F}(r_1, r_2, \rho) = \frac{1}{2} \frac{e^{\gamma_j(r_1 - r_2) - (\gamma_j^2/2)}}{r_1 r_2} G(r_1, r_2, \rho).$$

◇

6. Transition density of the Bessel process. In this section, we give some estimates for the transition density $(\tilde{q}_t(x, y))_{t>0}$, $x, y \geq 0$ of the three-dimensional Bessel process. As noted previously, we have

$$\tilde{q}_t(x, y) = \frac{1}{x} q_t(x, y) y, \quad x, y > 0, t > 0,$$

where $q_t(x, y)$ is the density of the Brownian motion killed at 0, which can be written

$$q_t(x, y) = g_t(x - y) - g_t(x + y),$$

where g_t is the density of the normal distribution with mean 0 and variance t . For $x = 0$, we have

$$\tilde{q}_t(0, y) = \sqrt{\frac{2}{\pi}} \frac{y^2}{t^{3/2}} e^{-y^2/(2t)}, \quad y > 0, t > 0.$$

We set, for $r > 0$, $m > 0$,

$$\bar{q}_t(r, m) = \frac{\tilde{q}_t(r, m)}{m} = \frac{1}{r} q_t(r, m).$$

Note that

$$\begin{aligned} \bar{q}_t(r, m) &= \frac{1}{r} (g_t(r - m) - g_t(r + m)) \\ &= - \int_{-1}^{+1} g'_t(m + r\xi) d\xi. \end{aligned}$$

This last expression allows to extend by continuity the definition of $\bar{q}_t(r, m)$ for $r = 0$ or $m = 0$, so that

$$\bar{q}_t(0, m) = - \int_{-1}^{+1} g'_t(m) d\xi = -2g'_t(m) = \sqrt{\frac{2}{\pi}} \frac{m}{t^{3/2}} e^{-m^2/(2t)}.$$

Notice that, for any $r \geq 0$, $\bar{q}_t(r, 0) = 0$.

PROPOSITION 6.1. *We have the following estimates, for any $\gamma \in \mathbb{R}$.*

1. *For any $s, r, m > 0$,*

$$\bar{q}_s(r, m) e^{\gamma m - (\gamma^2 s/2)} \leq \frac{1}{s} e^{\gamma + r} (C_1 + C_2 \gamma_+ \sqrt{s}),$$

with $C_1 = \sqrt{\frac{2}{\pi e}}$ and $C_2 = \sqrt{2/\pi}$.

2. For any $r, m > 0$,

$$\int_0^1 ds e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) \leq 2^{3/2} e^{r\gamma+}.$$

3. For any $s, r > 0$,

$$\int_0^{+\infty} dm e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) \leq \frac{4\sqrt{2}}{\sqrt{\pi s}} e^{r\gamma+}.$$

Proof. Note that we can assume that $\gamma \geq 0$ because, for $\gamma < 0$, $e^{\gamma m - (\gamma^2 s/2)} \leq 1$. We have, using the equalities $g_t(x) = g_1(x/\sqrt{t})/\sqrt{t}$ and $g_1'(x) = -xg_1(x)$

$$\begin{aligned} \bar{q}_s(r, m) &= - \int_{-1}^{+1} g_s'(m + r\xi) d\xi \\ &= \frac{1}{s} \int_{-1}^{+1} \frac{m + r\xi}{\sqrt{s}} g_1\left(\frac{m + r\xi}{\sqrt{s}}\right) d\xi. \end{aligned}$$

Note that

$$e^{\gamma m - (\gamma^2 s/2)} g_1\left(\frac{m + r\xi}{\sqrt{s}}\right) = e^{-\gamma r\xi} g_1\left(\frac{m + r\xi - \gamma s}{\sqrt{s}}\right).$$

Hence

$$\begin{aligned} e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) &= \frac{1}{s} \int_{-1}^{+1} e^{-\gamma r\xi} \frac{m + r\xi}{\sqrt{s}} g_1\left(\frac{m + r\xi - \gamma s}{\sqrt{s}}\right) d\xi \\ &\leq \frac{e^{r\gamma}}{s} \int_{-1}^{+1} \left| \frac{m + r\xi - \gamma s}{\sqrt{s}} + \gamma\sqrt{s} \right| g_1\left(\frac{m + r\xi - \gamma s}{\sqrt{s}}\right) d\xi \\ &\leq 2 \frac{e^{r\gamma}}{s} \left(\sup_{x>0} xg_1(x) + \gamma\sqrt{s} \frac{1}{\sqrt{2\pi}} \right), \end{aligned} \tag{6.1}$$

which gives the first inequality. For the second and third inequality, we start from (6.1) and notice that

$$\begin{aligned} (m + r\xi - \gamma s)^2 &= (m + r\xi)^2 + \gamma^2 s^2 - 2\gamma s(m + r\xi) \\ &\geq (m + r\xi)^2 + \gamma^2 s^2 - \left(2\gamma^2 s^2 + \frac{(m + r\xi)^2}{2} \right) \\ &= \frac{(m + r\xi)^2}{2} - \gamma^2 s^2. \end{aligned}$$

We deduce

$$g_1\left(\frac{m + r\xi - \gamma s}{\sqrt{s}}\right) \leq e^{\gamma^2 s/2} g_1\left(\frac{m + r\xi}{\sqrt{2s}}\right).$$

Hence

$$e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) \leq e^{\gamma r} \int_{-1}^{+1} \frac{|m + r\xi|}{s^{3/2}} g_1\left(\frac{m + r\xi}{\sqrt{2s}}\right) d\xi, \tag{6.2}$$

and, integrating with respect to s ,

$$\begin{aligned} \int_0^1 ds e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) &\leq e^{\gamma r} \int_{-1}^{+1} \left(\int_0^1 \frac{|m + r\xi|}{s^{3/2}} g_1\left(\frac{m + r\xi}{\sqrt{2s}}\right) ds \right) d\xi \\ &= 2^{3/2} e^{\gamma r} \int_{-1}^{+1} \left(\int_{|m+r\xi|/\sqrt{2}}^{+\infty} g_1(u) du \right) d\xi \\ &\leq 2^{3/2} e^{\gamma r}, \end{aligned}$$

where we have set $u = |m + r\xi|/\sqrt{2s}$. Integrating (6.2) with respect to m , we get

$$\begin{aligned} \int_0^{+\infty} e^{\gamma m - (\gamma^2 s/2)} \bar{q}_s(r, m) dm &\leq e^{\gamma r} \int_{-1}^{+1} \left(\int_0^{+\infty} \frac{|m + r\xi|}{s^{3/2}} g_1\left(\frac{m + r\xi}{\sqrt{2s}}\right) dm \right) d\xi \\ &\leq e^{\gamma r} \int_{-1}^{+1} \left(\int_{-\infty}^{+\infty} \frac{|m + r\xi|}{s^{3/2}} g_1\left(\frac{m + r\xi}{\sqrt{2s}}\right) dm \right) d\xi \\ &= 4 \frac{e^{\gamma r}}{s^{1/2}} \int_{-\infty}^{+\infty} |z| g_1(z) dz = 4\sqrt{2} \frac{e^{\gamma r}}{\sqrt{s\pi}}, \end{aligned}$$

where we have set $z = (m + r\xi)/\sqrt{2s}$. \diamond

We complete this section with a lemma concerning the minimum of the Bessel process. This result is a consequence of Lemma 3 of [1].

LEMMA 6.2. *Let $(R(t))_{t \geq 0}$ be a three-dimensional Bessel process starting from 0 and let t_1, t_2, y, m, b be positive numbers, with $t_1 < t_2$. We have, using the notation $R^\sharp(t_1, t_2) = \min_{u \in [t_1, t_2]} R(u)$,*

$$\mathbb{P}\left(R^\sharp(t_1, t_2) \leq b \mid R(t_1) = y, R(t_2) = m\right) \leq \frac{b(m+y)}{ym}.$$

Proof. We assume that $b < y \wedge m$, since if $b \geq y \wedge m$, the upper bound is larger than or equal to 1. We then have, using Lemma 3 of [1] (and the fact that the Bessel bridge can be viewed as a Brownian bridge conditioned to remain positive: see the proof of Lemma 4 of [1]),

$$\mathbb{P}\left(R^\sharp(t_1, t_2) \leq b \mid R(t_1) = y, R(t_2) = m\right) = \frac{e^{2(b-y)(m-b)/T} - e^{-2ym/T}}{1 - e^{-2ym/T}},$$

with $T = t_2 - t_1$. Hence, using the convexity of the exponential function and the inequality $b(m-b+y) \leq ym$,

$$\begin{aligned} \mathbb{P}\left(R^\sharp(t_1, t_2) \leq b \mid R(t_1) = y, R(t_2) = m\right) &= \frac{e^{2((b-y)(m-b)+ym)/T} - 1}{e^{2ym/T} - 1} \\ &= \frac{e^{2b(m-b+y)/T} - 1}{e^{2ym/T} - 1} \\ &\leq \frac{b(m-b+y)}{ym} \leq \frac{b(m+y)}{ym}. \end{aligned}$$

\diamond

7. Domination of the conditional probability. The following proposition will be used to ensure the domination of conditional expectations.

PROPOSITION 7.1. *There exists a constant $C_{t,\gamma,\sigma}$ (depending only on t, γ and σ) such that, for any $l \in \mathbb{N}$, $\theta = (0 < t_1 < \dots < t_l < t) \in \mathbb{R}^l$ et $\xi \in \mathbb{R}^l$, we have, for $j = 0, \dots, l$,*

$$\mathbb{P}_{l,\theta,\xi}(M^j \geq x > M^{j,n}) \leq \mathbb{1}_{\{t_{j+1}-t_j \leq 8t/n\}} + \frac{C_{t,\gamma,\sigma}}{\sqrt{n}} \frac{1}{\sqrt{t_{j+1}-t_j}}.$$

Proof. Note that $\mathbb{P}_{l,\theta,\xi}(M^j \geq x > M^{j,n}) = \mathbb{E}_{l,\theta,\xi} F(M^j, M^j - M^{j,n})$, if we define the function F by $F(m, \rho) = \mathbb{1}_{\{x \leq m < x+\rho\}}$. It follows from Proposition 5.2 that

$$\mathbb{P}_{l,\theta,\xi}(M^j \geq x > M^{j,n}) = \int_0^1 ds \mathbb{E} \left(L_s^j \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,n}\}} \right),$$

where L_s^j is given by (5.3) and

$$\tilde{x}_j = \frac{x - x_j}{\sigma_j}.$$

We can obviously assume that $t_{j+1} - t_j > 8t/n$, which, with the notations of Section 5, can be written $\lambda_j/n < 1/8$ and ensures that at least one of the sets I_n^- and I_n^+ is non-empty. So we can bound the random variable $R_s^{j,n}$ by $R^*(\lambda_j/n)$ where, for $u \in [0, 1]$, we set

$$R^*(u) = \max_{-u \leq v \leq +u} \check{R}(v).$$

Hence

$$\mathbb{P}_{l,\theta,\xi}(M^j \geq x > M^{j,n}) \leq \int_0^1 ds \mathbb{E} (L_s^j I_j(R_1(s), R^*(\lambda_j/n))),$$

where

$$I_j(r, \rho) = \mathbb{1}_{\{\tilde{x}_j \leq r \leq \tilde{x}_j + \rho\}}.$$

We have

$$\int_0^{\lambda_j/n} ds \mathbb{E} (L_s^j I_j(R_1(s), R^*(\lambda_j/n))) \leq \int_0^{\lambda_j/n} ds \mathbb{E} (L_s^j),$$

and

$$\mathbb{E} (L_s^j) = \mathbb{E} \left(\frac{e^{\gamma_j R_1(s) - \frac{\gamma_j^2 s}{2}}}{\sqrt{2} R_1(s)} \right) \mathbb{E} \left(\frac{e^{\gamma_j R_2(1-s) - \frac{\gamma_j^2 (1-s)}{2}}}{\sqrt{2} R_2(1-s)} \right).$$

By scaling, we have

$$\begin{aligned} \mathbb{E} \left(\frac{e^{\gamma_j R_1(s) - \frac{\gamma_j^2 s}{2}}}{\sqrt{2} R_1(s)} \right) &= \mathbb{E} \left(\frac{e^{\gamma_j \sqrt{s} R_1(1) - \frac{\gamma_j^2 s}{2}}}{\sqrt{2} \sqrt{s} R_1(1)} \right) \\ &\leq C_{t,\gamma,\sigma} \frac{1}{\sqrt{s}}. \end{aligned}$$

Similarly

$$\mathbb{E} \left(\frac{e^{\gamma_j R_2(1-s) - \frac{\gamma_j^2(1-s)}{2}}}{\sqrt{2} R_2(1-s)} \right) \leq C_{t,\gamma,\sigma} \frac{1}{\sqrt{1-s}}.$$

Therefore

$$\int_0^{\lambda_j/n} ds \mathbb{E} (L_s^j) \leq C_{t,\gamma,\sigma} \sqrt{\frac{\lambda_j}{n}} = \frac{C_{t,\gamma,\sigma}}{\sqrt{t_{j+1} - t_j}} \frac{1}{\sqrt{n}},$$

and, by a similar argument, $\int_{1-\lambda_j/n}^1 ds \mathbb{E} (L_s^j) \leq \frac{C_{t,\gamma,\sigma}}{\sqrt{t_{j+1} - t_j}} \frac{1}{\sqrt{n}}$. It remains to study the integral on the interval $[\lambda_j/n, 1 - \lambda_j/n]$. Denote by $(\mathcal{F}_s)_{s \geq 0}$ the natural filtration of the pair (R^1, R^2) . For $s \in [\lambda_j/n, 1 - \lambda_j/n]$, we have, by conditioning with respect to $\mathcal{F}_{\lambda_j/n}$,

$$\begin{aligned} & \mathbb{E} (L_s^j \mathbf{1}_{\{R_1(s) \in [\tilde{x}_j, \tilde{x}_j + R^*(\lambda_j/n)]\}}) \\ &= \mathbb{E} \left(\int_0^\infty dm \bar{q}_{s-\lambda_j/n}^j(R_1(\lambda_j/n), m) \phi_{1-s-\lambda_j/n}^j(R_2(\lambda_j/n)) I_j(m, R^*(\lambda_j/n)) \right), \end{aligned}$$

where

$$\bar{q}_s^j(r, m) = e^{\gamma_j m - \gamma_j^2/2} \bar{q}_s(r, m) \quad \text{and} \quad \phi_s^j(r) = \frac{1}{2} \int_0^\infty dm e^{-\gamma_j m} \bar{q}_s(r, m).$$

By Proposition 6.1, we have

$$\phi_s^j(r) \leq \frac{2\sqrt{2}}{\sqrt{\pi s}} e^{r|\gamma_j| + (\gamma_j^2/2)} \leq \frac{C_{t,\gamma,\sigma}}{\sqrt{s}} e^{r|\gamma_j|}.$$

We have

$$\begin{aligned} & \int_{\lambda_j/n}^{1-\lambda_j/n} ds \bar{q}_{s-\lambda_j/n}^j(R_1(\lambda_j/n), m) \phi_{1-s-\lambda_j/n}^j(R_2(\lambda_j/n)) \\ &= \int_0^{1-2\lambda_j/n} ds \bar{q}_s^j(R_1(\lambda_j/n), m) \phi_{1-s-2\frac{\lambda_j}{n}}^j(R_2(\lambda_j/n)). \end{aligned}$$

For $s \in [0, 1/2 - (\lambda_j/n)]$, we have

$$\frac{1}{\sqrt{1-s-2\lambda_j/n}} \leq \frac{1}{\sqrt{1/2 - (\lambda_j/n)}} \leq 2,$$

because $\lambda_j/n < 1/4$. Hence

$$\begin{aligned} & \int_{\lambda_j/n}^{1/2 - (\lambda_j/n)} ds \bar{q}_{s-\lambda_j/n}^j(R_1(\lambda_j/n), m) \phi_{1-s-\lambda_j/n}^j(R_2(\lambda_j/n)) \\ & \leq C_{t,\gamma,\sigma} \int_0^1 ds \bar{q}_s^j(R_1(\lambda_j/n), m) e^{|\gamma_j| R_2(\lambda_j/n)} \\ & \leq C_{t,\gamma,\sigma} e^{|\gamma_j|(R_1(\lambda_j/n) + R_2(\lambda_j/n))}, \end{aligned}$$

where the last inequality follows from Proposition 6.1. On the other hand, for $s \in [1/2 - (\lambda_j/n), 1 - \lambda_j/n]$, we have, using the first inequality of Proposition 6.1,

$$\bar{q}_{s-\lambda_j/n}^j(r, m) \leq \frac{C_{t,\gamma,\sigma}}{s - \lambda_j/n} e^{r|\gamma_j|} \leq 4C_{t,\gamma,\sigma} e^{r|\gamma_j|}.$$

Here, the last inequality follows from the condition $\lambda_j/n < 1/8$. We deduce

$$\begin{aligned} & \int_{1/2 - (\lambda_j/n)}^{1 - (\lambda_j/n)} ds \bar{q}_{s-\lambda_j/n}^j(R_1(\lambda_j/n), m) \phi_{1-s-\lambda_j/n}^j(R_2(\lambda_j/n)) \\ & \leq C_{t,\gamma,\sigma} e^{|\gamma_j| R_1(\lambda_j/n)} \int_{1/2}^1 ds \phi_{1-s}^j(R_2(\lambda_j/n)) \\ & \leq C_{t,\gamma,\sigma} e^{|\gamma_j|(R_1(\lambda_j/n) + R_2(\lambda_j/n))} \int_{1/2}^1 \frac{ds}{\sqrt{1-s}} \\ & \leq C_{t,\gamma,\sigma} e^{|\gamma_j|(R_1(\lambda_j/n) + R_2(\lambda_j/n))}. \end{aligned}$$

Then we have

$$\begin{aligned} & \int_{\lambda_j/n}^{1-\lambda_j/n} ds \mathbb{E} \left(L_s^j \mathbf{1}_{\{R_1(s) \in [\bar{x}_j, \bar{x}_j + R^*(\lambda_j/n)]\}} \right) \\ & \leq C_{t,\gamma,\sigma} \mathbb{E} \left(e^{|\gamma_j|(R_1(\lambda_j/n) + R_2(\lambda_j/n))} \int_0^\infty dm I_j(m, R^*(\lambda_j/n)) \right) \quad (7.1) \\ & \leq C_{t,\gamma,\sigma} \mathbb{E} \left(R^*(\lambda_j/n) e^{|\gamma_j|(R_1(\lambda_j/n) + R_2(\lambda_j/n))} \right) \\ & = C_{t,\gamma,\sigma} \sqrt{\lambda_j/n} \mathbb{E} \left(R^*(1) e^{|\gamma_j| \sqrt{\lambda_j/n} (R_1(1) + R_2(2))} \right) \\ & = \frac{C_{t,\gamma,\sigma}}{\sqrt{t_{j+1} - t_j}} \frac{1}{\sqrt{n}}, \end{aligned}$$

where we have used the scaling property of the Bessel process, $\sqrt{\lambda_j} = \sqrt{t/(t_{j+1} - t_j)}$, and $|\gamma_j| \sqrt{\lambda_j} = |\gamma| \sqrt{t}/\sigma$. \diamond

REMARK 7.2. It follows from the proof of the proposition that, for any $\delta > 0$, we have

$$\mathbb{P}_{l,\theta,\xi} (M^j \geq x > M^j - \delta/\sqrt{n}) \leq \mathbf{1}_{\{t_{j+1} - t_j \leq 8t/n\}} + \frac{C_{t,\gamma,\sigma,\delta}}{\sqrt{n}} \frac{1}{\sqrt{t_{j+1} - t_j}}.$$

Indeed, we have $\mathbb{P}_{l,\theta,\xi} (M^j \geq x > M^j - \delta/\sqrt{n}) = \int_0^1 ds \mathbb{E} (L_s^j I_j(R_1(s), \delta/(\sigma_j \sqrt{n})))$, and we can replace $R^*(\lambda_j/n)$ with $\delta/(\sigma_j \sqrt{n})$ in (7.1).

8. Convergence of the conditional expectation. The aim of this section is to prove the following result and to deduce Theorem 2.1.

THEOREM 8.1. *Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a bounded Borel measurable function, such that $m \mapsto F(y, m)$ is continuous for all $y \in \mathbb{R}$.*

We have, with the notation $\mathbb{E}_0 = \mathbb{E}(\cdot \mid N_t = 0)$,

$$\mathbb{E}_0 (F(X_t, M^0) \mathbf{1}_{\{M^0 \geq x > M^0, n\}}) = \mathbb{E}_0 \left(F(X_t, M^0) \mathbf{1}_{\{M^0 \geq x > M^0 - \sigma_{\beta_1} \sqrt{t/n}\}} \right) + o(1/\sqrt{n}).$$

Moreover, for any positive integer l , and for any $\theta = (t_1, \dots, t_l) \in \mathbb{R}^l$, with $0 < t_1 < \dots < t_l < t$, $\xi = (x_1, \dots, x_l) \in \mathbb{R}^l$, we have, for $j \in \{0, 1, \dots, l\}$, if $x_j \neq x$,

$$\mathbb{E}_{l,\theta,\xi} \left(F(X_{t_{j+1}^-}, M^j) \mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \right) = \mathbb{E}_{l,\theta,\xi} \left(F(X_{t_{j+1}^-}, M^j) \mathbb{1}_{\{M^j \geq x > M^j - \sigma \beta_1 \sqrt{t/n}\}} \right) + o(1/\sqrt{n}),$$

where β_1 is defined as in Theorem 2.1.

We will first show how Theorem 2.1 can be deduced from Theorem 8.1.

Proof of Theorem 2.1. Observe that, with the notation $\mathbb{E}_0 = \mathbb{E}(\cdot \mid N_t = 0)$,

$$\begin{aligned} \mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) &= \mathbb{E}_0 \left(g(X_t) \mathbb{1}_{\{M^0 \geq x > M^{0,n}\}} \right) \mathbb{P}(N_t = 0) \\ &\quad + \mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \mid N_t \geq 1 \right) \mathbb{P}(N_t \geq 1). \end{aligned}$$

Using Theorem 8.1, we have

$$\mathbb{E}_0 \left(g(X_t) \mathbb{1}_{\{M^0 \geq x > M^{0,n}\}} \right) = \mathbb{E}_0 \left(g(X_t) \mathbb{1}_{\{M^0 \geq x > M^0 - \sigma \beta_1 \sqrt{t/n}\}} \right) + o(1/\sqrt{n}).$$

On the other hand, we deduce from Proposition 4.1 that

$$\mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \mid N_t \geq 1 \right) = \mathbb{E} \left(\sum_{j=0}^{N_t} \alpha_{N_t}^{j,n}(T_1, \dots, T_n) \mid N_t \geq 1 \right) + o(1/\sqrt{n}),$$

where, for any positive integer l , and for $j = 0, \dots, l$,

$$\alpha_l^{j,n}(\theta) = \mathbb{E}_{l,\theta} \left(g(X_t) \mathbb{1}_{\{M^j \geq x > M^{j,n}, M^j > \max_{i \neq j} M^i\}} \right).$$

Note that, as a consequence of Lemma 5.1, we have, assuming $x_j \neq x$,

$$\begin{aligned} \mathbb{E}_{l,\theta,\xi} \alpha_l^{j,n}(\theta) &= \mathbb{E}_{l,\theta,\xi} \left(\mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \alpha_{l,\theta,\xi}^j(X_{t_{j+1}^-}, M^j) \right) \\ &= \mathbb{E}_{l,\theta,\xi} \left(\mathbb{1}_{\{M^j \geq x > M^j - \beta_1 \sigma \sqrt{t/n}\}} \alpha_{l,\theta,\xi}^j(X_{t_{j+1}^-}, M^j) \right) + o(1/\sqrt{n}) \\ &= \mathbb{E}_{l,\theta,\xi} \left(\mathbb{1}_{\{M^j \geq x > M^j - \beta_1 \sigma \sqrt{t/n}, M^j > \max_{i \neq j} M^i\}} \right) + o(1/\sqrt{n}), \end{aligned}$$

where the second equality follows from Theorem 8.1, and the last one from the expression of $\alpha_{l,\theta,\xi}^j$ (see Lemma 5.1 and its proof; note that $m \mapsto \alpha_{l,\theta,\xi}^j(y, m)$ is continuous because $\mathbb{P}_{l,\theta,\xi}(\max_{i \neq j} M_i = m) = 0$). By taking the sum over $j = 0, \dots, l$, we deduce that, for l , θ and ξ fixed, we have

$$\mathbb{E}_{l,\theta,\xi} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) = \mathbb{E}_{l,\theta,\xi} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t - \beta_1 \sigma \sqrt{t/n}\}} \right) + o(1/\sqrt{n}). \quad (8.1)$$

Observe that $\mathbb{P}(X_{T_j} = x) = 0$ for all jump times T_j (including $T_0 = 0$, since $x > 0$). Therefore, in order to get the result for the unconditional expectations, we only need to check a domination condition. We have

$$\begin{aligned} \left| \mathbb{E}_{l,\theta,\xi} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) \right| &\leq \|g\|_\infty \mathbb{P}_{l,\theta,\xi}(M_t \geq x > M_t^n) \\ &\leq \|g\|_\infty \sum_{j=0}^l \mathbb{P}_{l,\theta,\xi}(M^j \geq x > M^{j,n}), \end{aligned}$$

and

$$\mathbb{P}_{l,\theta,\xi} \left(M_t \geq x > M_t - \beta_1 \sigma \sqrt{t/n} \right) \leq \sum_{j=0}^l \mathbb{P}_{l,\theta,\xi} \left(M^j \geq x > M^j - \beta_1 \sigma \sqrt{t/n} \right).$$

Using Proposition 7.1 and Remark 7.2, we deduce that

$$\mathbb{P}_{l,\theta,\xi} (M_t \geq x > M_t^n) + \mathbb{P}_{l,\theta,\xi} (M_t \geq x > M_t - \beta_1 \sigma \sqrt{t/n}) \leq P(l, \theta),$$

where

$$P(l, \theta) = P_l(t_1, \dots, t_l) = \sum_{j=0}^l \mathbb{1}_{\{t_{j+1}-t_j \leq 8t/n\}} + \frac{C_{t,\gamma,\sigma}}{\sqrt{n}} \sum_{j=0}^l \frac{1}{\sqrt{t_{j+1}-t_j}}$$

It follows from Proposition 3.2 that

$$\mathbb{E} \left(\sum_{j=0}^{N_t} \mathbb{1}_{\{T_{j+1} \wedge t - T_j \leq 8t/n\}} \right) \leq \frac{8t}{n} \mathbb{E} (N_t(N_t + 1)),$$

and, from Proposition 3.1, that

$$\mathbb{E} \left(\sum_{j=0}^{N_t} \frac{1}{\sqrt{T_{j+1} \wedge t - T_j}} \right) \leq 2 \frac{\mathbb{E} (N_t(N_t + 1))}{\sqrt{t}}.$$

The last two inequalities are sufficient to extend the estimate (8.1) to unconditional expectations. \diamond

For the proof of Theorem 8.1, we start from the representation given by Proposition 5.2, which reads

$$\mathbb{E}_{l,\theta,\xi} \left(F(X_{t_{j+1}^-}, M^j) \mathbb{1}_{\{M^j \geq x > M^{j,n}\}} \right) = \int_0^1 E_{l,\theta,\xi}^{j,n}(s) ds,$$

with

$$E_{l,\theta,\xi}^{j,n}(s) = \mathbb{E} \left(L_s^j \alpha_j(R_1(s), R_2(1-s)) \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,n}\}} \right),$$

where $L_s^j = \frac{e^{\gamma_j(R_1(s) - R_2(1-s)) - (\gamma_j^2/2)}}{2R_1(s)R_2(1-s)}$,

$$\alpha_j(r_1, r_2) = F(x_j + \sigma_j(r_1 - r_2), x_j + \sigma_j r_1), \text{ and } \tilde{x}_j = \frac{x - x_j}{\sigma_j}.$$

Note that the function α_j is bounded, and $\|\alpha_j\|_\infty = \|F\|_\infty$. For any integer $J \geq 1$, we can write, for n large enough,

$$\int_0^1 E_{l,\theta,\xi}^{j,n}(s) ds = \int_0^{\frac{\lambda_j(J+1)}{n}} E_{l,\theta,\xi}^{j,n}(s) ds + \int_{1-\frac{\lambda_j J}{n}}^1 E_{l,\theta,\xi}^{j,n}(s) ds + \int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} E_{l,\theta,\xi}^{j,n}(s) ds. \quad (8.2)$$

The first two terms of this decomposition are controlled via the following lemma.

LEMMA 8.2. *For any integer $J \geq 1$, for any (l, θ, ξ) , and for $j = 0, 1, \dots, l$, we have, if $x_j \neq 0$*

$$\int_0^{\frac{\lambda_j(J+1)}{n}} |E_{l,\theta,\xi}^{j,n}(s)| ds + \int_{1-\frac{\lambda_j J}{n}}^1 |E_{l,\theta,\xi}^{j,n}(s)| ds = o(1/\sqrt{n}).$$

Proof. We will only consider the first integral, the argument is similar for the second term. Note that for n large enough and $s \in [0, \lambda_j(J+1)/n]$, we have $1-s > \lambda_j/n$ and we can bound $R_s^{j,n}$ from above by $R_2^*(\lambda_j/n)$ (with $R_2^*(s) = \max_{0 \leq u \leq s} R_2^*(u)$). Then we have, using the boundedness of F ,

$$\left| E_{l,\theta,\xi}^{j,n}(s) \right| \leq \|F\|_\infty \mathbb{E} \left(\tilde{L}_s^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*\left(\frac{\lambda_j}{n}\right)\}} \right),$$

where $\tilde{L}_s^j = \frac{e^{|\gamma_j|(R_1(s)+R_2(1-s))}}{2R_1(s)R_2(1-s)}$. Hence, with the substitution $s' = ns$,

$$\begin{aligned} \sqrt{n} \int_0^{\lambda_j(J+1)/n} \left| E_{l,\theta,\xi}^{j,n}(s) \right| ds &\leq \sqrt{n} \|F\|_\infty \int_0^{\lambda_j(J+1)/n} ds \mathbb{E} \left(\tilde{L}_s^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*\left(\frac{\lambda_j}{n}\right)\}} \right) \\ &= \|F\|_\infty \int_0^{\lambda_j(J+1)} \frac{ds'}{\sqrt{n}} \mathbb{E} \left(\tilde{L}_{s'/n}^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s'/n) \leq \tilde{x}_j + R_2^*\left(\frac{\lambda_j}{n}\right)\}} \right). \end{aligned}$$

By scaling, we can write, using the notation

$$\Lambda_j(r_1, r_2, \rho) = \frac{e^{|\gamma_j|(r_1+r_2)}}{2r_1r_2} \mathbf{1}_{\{\tilde{x}_j \leq r_1 \leq \tilde{x}_j + \rho\}},$$

$$\begin{aligned} \mathbb{E} \left(\tilde{L}_{s/n}^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s/n) \leq \tilde{x}_j + R_2^*\left(\frac{\lambda_j}{n}\right)\}} \right) &= \mathbb{E} \left(\Lambda_j \left(R_1(s/n), R_2(1 - \frac{s}{n}), R_2^*\left(\frac{\lambda_j}{n}\right) \right) \right) \\ &= \mathbb{E} \left(\Lambda_j \left(R_1(s)/\sqrt{n}, R_2(1 - \frac{s}{n}), R_2^*\left(\frac{\lambda_j}{n}\right) \right) \right) \\ &\leq \sqrt{n} \mathbb{E} \left(\Lambda_j^{s,n} \left(R_1(s), R_2(1), R_2^*\left(\frac{\lambda_j}{n-s}\right) \right) \right), \end{aligned}$$

with

$$\Lambda_j^{s,n}(r_1, r_2, \rho) = \frac{e^{|\gamma_j|\left(\frac{r_1}{\sqrt{n}} + \sqrt{1-\frac{s}{n}}r_2\right)}}{2r_1\sqrt{1-\frac{s}{n}}r_2} \mathbf{1}_{\{\tilde{x}_j \leq \frac{r_1}{\sqrt{n}} \leq \tilde{x}_j + \rho\}}.$$

Hence

$$\sqrt{n} \int_0^{\lambda_j(J+1)/n} \left| E_{l,\theta,\xi}^{j,n}(s) \right| ds \leq \|F\|_\infty \int_0^{\lambda_j(J+1)} ds \mathbb{E} \left(\Lambda_j^{s,n} \left(R_1(s), R_2(1), R_2^*\left(\frac{\lambda_j}{n-s}\right) \right) \right).$$

Now, if $n > 2\lambda_j(J+1)$, we have, for $s \in [0, \lambda_j(J+1)]$, $n-s \geq n/2$, so that $R_2^*\left(\frac{\lambda_j}{n-s}\right) \leq R_2^*\left(\frac{2\lambda_j}{n}\right)$, and $\sqrt{1-s/n} \geq 1/\sqrt{2}$, so that

$$\Lambda_j^{s,n} \left(R_1(s), R_2(1), R_2^*\left(\frac{\lambda_j}{n-s}\right) \right) \leq \frac{e^{|\gamma_j|(R_1(s)+R_2(1))}}{\sqrt{2}R_1(s)R_2(1)} \mathbf{1}_{\{\tilde{x}_j \leq R_1(s)/\sqrt{n} \leq \tilde{x}_j + R_2^*\left(\frac{2\lambda_j}{n}\right)\}}.$$

If $x \neq x_j$, we have $\tilde{x}_j \neq 0$, and the right hand side of the inequality goes to 0 almost surely as $n \rightarrow \infty$, for all $s \in (0, 1)$. Since $\int_0^1 ds \mathbb{E} \left(\frac{e^{|\gamma_j|(R_1(s)+R_2(1))}}{\sqrt{2}R_1(s)R_2(1)} \right) < \infty$, we conclude that

$$\lim_{n \rightarrow \infty} \left(\sqrt{n} \int_0^{\lambda_j(J+1)/n} E_{l,\theta,\xi}^{j,n}(s) ds \right) = 0.$$

◇

We will now examine the case $x < x_j$.

LEMMA 8.3. *If $x < x_j$, we have,*

$$\int_0^1 |E_{l,\theta,\xi}^{j,n}(s)| ds = o(1/\sqrt{n}).$$

Proof. In view of Lemma 8.2, it suffices to show that $\int_{\lambda_j/n}^{1-\lambda_j/n} |E_{l,\theta,\xi}^{j,n}(s)| ds = o(1/\sqrt{n})$. From (7.1), we have, for n large enough,

$$\begin{aligned} \int_{\lambda_j/n}^{1-\lambda_j/n} |E_{l,\theta,\xi}^{j,n}(s)| ds &\leq C \mathbb{E} \left(e^{|\gamma_j|(R_1(\lambda_j/n)+R_2(\lambda_j/n))} \int_0^\infty dm \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + R^*(\lambda_j/n)\}} \right) \\ &\leq C \mathbb{E} \left(e^{|\gamma_j|(R_1(\lambda_j/n)+R_2(\lambda_j/n))} R^*(\lambda_j/n) \mathbb{1}_{\{R^*(\lambda_j/n) \geq -\tilde{x}_j\}} \right) \\ &= C \sqrt{\frac{\lambda_j}{n}} \mathbb{E} \left(e^{|\gamma_j| \sqrt{\frac{\lambda_j}{n}}(R_1(1)+R_2(1))} R^*(1) \mathbb{1}_{\{R^*(1) \geq -\tilde{x}_j \sqrt{n/\lambda_j}\}} \right) \end{aligned}$$

Since $\tilde{x}_j < 0$, the right hand side of the last equality is $o(1/\sqrt{n})$. ◇

We will now study the case $x_j < x$. We go back to the decomposition (8.2) and assume that n is large enough, so that $\lambda_j J/n < 1/4$. Note that, for $s \in [\lambda_j(J+1)/n, 1-\lambda_j J/n]$, we have $N_n^1 \geq J$ and $N_n^2 \geq J$. So, we have

$$R_s^{j,n} \leq R_s^{j,J,n},$$

where

$$R_s^{j,J,n} = \min_{-J \leq k \leq J} \check{R}(d_n^j(s) + \lambda_j(k/n)).$$

LEMMA 8.4. *If $x_j < x$, we have,*

$$\lim_{J \rightarrow +\infty} \limsup_{n \rightarrow \infty} \sqrt{n} \left(\int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} (E_{l,\theta,\xi}^{j,J,n}(s) - E_{l,\theta,\xi}^{j,n}(s)) ds \right) = 0,$$

where

$$E_{l,\theta,\xi}^{j,J,n}(s) = \mathbb{E} \left(L_s^j \alpha_j(R_1(s), R_2(1-s)) \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,J,n}\}} \right).$$

Proof. Note that

$$E_{l,\theta,\xi}^{j,J,n}(s) - E_{l,\theta,\xi}^{j,n}(s) = \mathbb{E} \left(L_s^j \alpha_j(R_1(s), R_2(1-s)) \mathbb{1}_{\Delta_s^{j,J,n}} \right),$$

with

$$\Delta_s^{j,J,n} = \left\{ \tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,J,n} \text{ and } \exists k \in [0, N_n^1] \cup [-N_n^2, -1], \right. \\ \left. \tilde{x}_j + \check{R}(d_n^j(s) + \lambda_j(k/n)) < R_1(s) \right\}.$$

Introducing the notation, for $i = 1, 2$, and for any real numbers s_1, s_2 with $0 < s_1 < s_2$,

$$R_i^\#(s_1, s_2) = \min_{u \in [s_1, s_2]} R_i(u),$$

we see that

$$\Delta_s^{j,J,n} \subset \left\{ \tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,J,n} \right\} \\ \cap \left\{ R_1^\#(\lambda_j J/n, s) < R_s^{j,J,n} \text{ or } R_2^\#(\lambda_j J/n, 1-s) < R_s^{j,J,n} \right\}.$$

Note that $R_s^{j,J,n} \leq R_1^*(\lambda_j/n) \wedge R_2^*(\lambda_j/n)$ and $R_2^\#(s_1, s_2) \leq R_2^\#(s_1)$, where

$$R_2^\#(s) = \min_{u \in [s, +\infty[} R_2^\#(u), \quad s \geq 0.$$

So, we have

$$|E_{l,\theta,\xi}^{j,J,n}(s) - E_{l,\theta,\xi}^{j,n}(s)| \leq \|F\|_\infty \left(F_{l,\theta,\xi}^{j,J,n}(s) + G_{l,\theta,\xi}^{j,J,n}(s) \right),$$

where

$$F_{l,\theta,\xi}^{j,J,n}(s) = \mathbb{E} \left(L_s^j \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_1^*(\lambda_j/n), R_2^\#(\lambda_j J/n) < R_1^*(\lambda_j/n)\}} \right)$$

and

$$G_{l,\theta,\xi}^{j,J,n}(s) = \mathbb{E} \left(L_s^j \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*(\lambda_j/n), R_1^\#(\lambda_j J/n, s) < R_2^*(\lambda_j/n)\}} \right).$$

In the sequel we denote by $(\mathcal{F}_s^i)_{s \geq 0}$ ($i = 1, 2$) the natural filtration of the process $(R_i(s))_{s \geq 0}$ and by \mathcal{F}^i the σ -algebra generated by the union of the σ -algebras \mathcal{F}_s^i , $s \geq 0$.

In order to estimate $F_{l,\theta,\xi}^{j,J,n}(s)$, we write $L_s^j = \frac{e^{\gamma_j R_1(s) - \frac{\gamma_j^2}{2}(s)}}{R_1(s)} M_{1-s}^j$, with

$$M_{1-s}^j = \frac{e^{-\gamma_j R_2(1-s) - \frac{\gamma_j^2}{2}(1-s)}}{2R_2(1-s)}.$$

By conditioning with respect to \mathcal{F}^1 and using the Cauchy-Schwarz inequality, we get

$$\mathbb{E} \left(\frac{e^{-\gamma_j R_2(1-s) - \frac{\gamma_j^2}{2}(1-s)}}{2R_2(1-s)} \mathbb{1}_{\{R_2^\#(\lambda_j J/n) < R_1^*(\lambda_j/n)\}} \mid \mathcal{F}^1 \right) \\ \leq \|M_{1-s}^j\|_2 \left(\mathbb{E} \left(\mathbb{1}_{\{R_2^\#(\lambda_j J/n) < R_1^*(\lambda_j/n)\}} \mid \mathcal{F}^1 \right) \right)^{1/2} \\ = \|M_{1-s}^j\|_2 \sqrt{f_J^{j,n}(R_1^*(\lambda_j/n))},$$

where $f_J^{j,n}$ is the cumulative distribution function of the random variable $R_2^\#(\lambda_j J/n)$. Note that, by scaling, we have, for any $r > 0$,

$$f_J^{j,n}(r) = f_J\left(r\sqrt{n/\lambda_j}\right), \quad (8.3)$$

where f_J denotes the cumulative distribution function of $R_2^\#(J)$. On the other hand, we have

$$\begin{aligned} \|M_{1-s}^j\|_2 &= \frac{1}{2\sqrt{1-s}} \left\| \frac{e^{-\gamma_j \sqrt{1-s} R_2(1) - \frac{\gamma_j^2}{2}(1-s)}}{R_2(1)} \right\|_2 \\ &= \frac{1}{2\sqrt{1-s}} \left(\int_0^\infty e^{-2\gamma_j \sqrt{1-s} m - \gamma_j^2(1-s) - \frac{m^2}{2}} \sqrt{\frac{2}{\pi}} dm \right)^{1/2} \\ &\leq \frac{1}{2\sqrt{1-s}} \left(\int_{-\infty}^\infty e^{-2\gamma_j \sqrt{1-s} m - \gamma_j^2(1-s) - \frac{m^2}{2}} \sqrt{\frac{2}{\pi}} dm \right)^{1/2} \\ &= \frac{1}{2\sqrt{1-s}} \left(2e^{\gamma_j^2(1-s)} \right)^{1/2} = \frac{1}{\sqrt{2(1-s)}} e^{\gamma_j^2(1-s)/2}. \end{aligned} \quad (8.4)$$

Using the inequalities (8.3) and (8.4) in the expression of $F_{l,\theta,\xi}^{j,J,n}(s)$, we get, after conditioning with respect to $\mathcal{F}_{\lambda_j/n}^1$,

$$\begin{aligned} F_{l,\theta,\xi}^{j,J,n}(s) &\leq \frac{e^{\gamma_j^2/2}}{\sqrt{2(1-s)}} \mathbb{E} \left(\sqrt{f_J^{j,n}(R_1^*(\lambda_j/n))} \right. \\ &\quad \times \left. \int_0^\infty \bar{q}_{s-\lambda_j/n}^j(R_1(\lambda_j/n), m) \mathbb{1}_{\{\bar{x}_j \leq m \leq \bar{x}_j + R_1^*(\lambda_j/n)\}} dm \right), \end{aligned}$$

with

$$\bar{q}_{s-\lambda_j/n}^j(r, m) = e^{\gamma_j m} \bar{q}_{s-\lambda_j/n}(r, m).$$

We can prove (by the same arguments as in the proof of Proposition 7.1) that

$$\int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} \bar{q}_{s-\lambda_j/n}^j(r, m) \frac{ds}{\sqrt{2(1-s)}} \leq C_{t,\gamma,\sigma} e^{r|\gamma_j|}.$$

Hence

$$\begin{aligned} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} F_{l,\theta,\xi}^{j,J,n}(s) ds &\leq C_{t,\gamma,\sigma} \mathbb{E} \left(\sqrt{f_J^{j,n}(R_1^*(\lambda_j/n))} e^{|\gamma_j| R_1(\lambda_j/n)} R_1^*(\lambda_j/n) \right) \\ &= C_{t,\gamma,\sigma} \mathbb{E} \left(\sqrt{f_J(R_1^*(1))} e^{|\gamma_j| \sqrt{\lambda_j/n} R_1(1)} \sqrt{\lambda_j/n} R_1^*(1) \right), \end{aligned}$$

where the last inequality follows from (8.3) and the scaling invariance of R_1 . Hence

$$\limsup_{n \rightarrow \infty} \sqrt{n} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} F_{l,\theta,\xi}^{j,J,n}(s) ds \leq C_{t,\gamma,\sigma} \sqrt{\lambda_j} \mathbb{E} \left(\sqrt{f_J(R_1^*(1))} R_1^*(1) \right).$$

We have $f_J(r) = \mathbb{P}(R_2^\sharp(J) \leq r) = \mathbb{P}(R_2^\sharp(1) \leq r/\sqrt{J})$. So, for any $r > 0$, $\lim_{J \rightarrow \infty} f_J(r) = 0$. We deduce, by dominated convergence that

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \sqrt{n} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} F_{l,\theta,\xi}^{j,J,n}(s) ds = 0.$$

We will now prove the same property for $G_{l,\theta,\xi}^{j,J,n}(s)$. We have, by conditioning with respect to the σ -algebra generated by \mathcal{F}^2 and the pair $(R_1(\lambda_j J/n), R_1(s))$,

$$\begin{aligned} G_{l,\theta,\xi}^{j,J,n}(s) &= \mathbb{E} \left(L_s^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*(\lambda_j/n), R_1^\sharp(\lambda_j J/n, s) < R_2^*(\lambda_j/n)\}} \right) \\ &= \mathbb{E} \left(L_s^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*(\lambda_j/n)\}} \Delta \left(\frac{\lambda_j J}{n}, s, R_1 \left(\frac{\lambda_j J}{n} \right), R_1(s), R_2^* \left(\frac{\lambda_j}{n} \right) \right) \right), \end{aligned}$$

with the notation, for $0 < t_1 < t_2$ and $b, y, m > 0$,

$$\Delta(t_1, t_2, y, m, b) = \mathbb{P} \left(R_1^\sharp(t_1, t_2) \leq b \mid R_1(t_1) = y, R_1(t_2) = m \right).$$

By Lemma 6.2, we have $\Delta(t_1, t_2, y, m, b) \leq \bar{\Delta}(y, m, b)$, with

$$\bar{\Delta}(y, m, b) = \frac{b(m+y)}{ym} \wedge 1.$$

Therefore

$$G_{l,\theta,\xi}^{j,J,n}(s) \leq \mathbb{E} \left(L_s^j \mathbf{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_2^*(\lambda_j/n)\}} \bar{\Delta}(R_1(\lambda_j J/n), R_1(s), R_2^*(\lambda_j/n)) \right).$$

Note that

$$\begin{aligned} \mathbb{E} \left(\frac{e^{-\gamma_j R_2(1-s) - \frac{\gamma_j^2}{2}(1-s)}}{2R_2(1-s)} \mid \mathcal{F}_{\lambda_j/n}^2 \vee \mathcal{F}^1 \right) &= \frac{1}{2} \int_0^\infty dm e^{-\gamma_j m - \frac{\gamma_j^2}{2}(1-s)} \\ &\quad \times \bar{q}_{1-s-\lambda_j/n}(R_2(\lambda_j/n), m) \\ &\leq \frac{C}{\sqrt{1-s-\lambda_j/n}} e^{|\gamma_j| R_2(\lambda_j/n)}, \end{aligned}$$

where we have used the third estimate of Proposition 6.1. Now, condition with respect to $\mathcal{F}_{\lambda_j J/n}^1 \vee \mathcal{F}^2$ to get (introducing the random interval $I_n^{*j} = [\tilde{x}_j, \tilde{x}_j + R_2^*(\lambda_j/n)]$ in notations)

$$\begin{aligned} G_{l,\theta,\xi}^{j,J,n}(s) &\leq C \mathbb{E} \left(\frac{e^{|\gamma_j|(R_2(\lambda_j/n) + R_1(s))}}{\sqrt{1-s-\lambda_j/n} R_1(s)} \mathbf{1}_{I_n^{*j}}(R_1(s)) \right. \\ &\quad \times \bar{\Delta}(R_1(\lambda_j J/n), R_1(s), R_2^*(\lambda_j/n)) \Big) \\ &= C \mathbb{E} \left(\frac{e^{|\gamma_j| R_2(\lambda_j/n)}}{\sqrt{1-s-\lambda_j/n}} \int_0^\infty dm \bar{q}_{s-\lambda_j J/n}(R_1(\lambda_j J/n), m) \right. \\ &\quad \times \bar{\Delta}_n^j(R_1(\lambda_j J/n), m, R_2^*(\lambda_j/n)) \Big), \end{aligned}$$

where we have set

$$\bar{\Delta}_n^j(r_1, m, r_2) = e^{|\gamma_j| m} \mathbf{1}_{I_n^{*j}}(m) \bar{\Delta}(r_1, m, r_2).$$

Note that (with the arguments of the proof of Proposition 7.1)

$$\int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} e^{|\gamma_j|m} \bar{q}_{s-\lambda_j J/n}(r, m) \frac{ds}{\sqrt{1-s-\lambda_j/n}} \leq C_{t,\gamma,\sigma} e^{r|\gamma_j|},$$

so that

$$\begin{aligned} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} ds G_{l,\theta,\xi}^{j,J,n}(s) &\leq C_{t,\gamma,\sigma} \mathbb{E} \left(e^{|\gamma_j|(R_2(\lambda_j/n)+R_1(\lambda_j J/n))} \int_0^\infty dm \mathbb{1}_{I_n^{*j}}(m) \right. \\ &\quad \times \bar{\Delta}(R_1(\lambda_j J/n), m, R_2^*(\lambda_j/n)) \\ &= C_{t,\gamma,\sigma} \mathbb{E} \left(e^{|\gamma_j|\sqrt{\lambda_j/n}(R_1(J)+R_2(1))} \int_0^\infty dm \mathbb{1}_{\tilde{I}_n^{*j}}(m) \right. \\ &\quad \times \tilde{\Delta}_n^j(R_1(J), m, R_2^*(1)) \Big), \end{aligned}$$

with

$$\tilde{I}_n^{*j} = \left[\tilde{x}_j, \tilde{x}_j + \sqrt{\lambda_j/n} R_2^*(1) \right], \quad \tilde{\Delta}_n^j(r_1, m, r_2) = \bar{\Delta} \left(\sqrt{\lambda_j/n} r_1, m, \sqrt{\lambda_j/n} r_2 \right).$$

We have

$$\begin{aligned} \tilde{\Delta}_n^j(R_1(J), m, R_2^*(1)) &= \left[R_2^*(1) \left(\frac{1}{R_1(J)} + \frac{\sqrt{\lambda_j/n}}{m} \right) \right] \wedge 1 \\ &\leq \frac{R_2^*(1)}{R_1(J)} + \left(R_2^*(1) \frac{\sqrt{\lambda_j/n}}{m} \right) \wedge 1. \end{aligned}$$

By assumption, $x_j < x$, so that $\tilde{x}_j > 0$. Therefore, for $m \in I_n^{*j}$, we have $1/m \leq 1/\tilde{x}_j$, so that

$$\int_0^\infty dm \mathbb{1}_{\tilde{I}_n^{*j}}(m) \tilde{\Delta}_n^j(R_1(J), m, R_2^*(1)) \leq \sqrt{\lambda_j/n} R_2^*(1) \left(\frac{R_2^*(1)}{R_1(J)} + R_2^*(1) \frac{\sqrt{\lambda_j/n}}{\tilde{x}_j} \right).$$

Hence

$$\limsup_{n \rightarrow \infty} \sqrt{n} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} ds G_{l,\theta,\xi}^{j,J,n}(s) \leq C_{t,\gamma,\sigma} \sqrt{\lambda_j} \mathbb{E} \left(\frac{R_2^*(1)^2}{R_1(J)} \right).$$

Since $\lim_{J \rightarrow \infty} R_1(J) = 0$, we have

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \sqrt{n} \int_{\lambda_j(J+1)/n}^{1-\lambda_j J/n} ds G_{l,\theta,\xi}^{j,J,n}(s) = 0.$$

◇

We will now study the asymptotic behavior of $\int_{\frac{\lambda_j J}{n}}^{1-\frac{\lambda_j J}{n}} E_{l,\theta,\xi}^{j,J,n}(s) ds$. Note that, by conditioning with respect to $\mathcal{F}_{\lambda_j J/n}^1 \vee \mathcal{F}_{\lambda_j J/n}^2$, we have

$$\begin{aligned} E_{l,\theta,\xi}^{j,J,n}(s) &= \mathbb{E} \left(L_s^j \alpha_j(R_1(s), R_2(1-s)) \mathbb{1}_{\{\tilde{x}_j \leq R_1(s) \leq \tilde{x}_j + R_s^{j,J,n}\}} \right) \\ &= \mathbb{E} \int_0^\infty dm \bar{q}_{s-\lambda_j J/n}^j(R_1(\lambda_j J/n), m) \bar{\phi}_{1-s-\lambda_j J/n}^j(R_2(\lambda_j J/n), m) \\ &\quad \times \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + R_s^{j,J,n}\}}, \end{aligned}$$

with the notations

$$\bar{q}_s^j(r, m) = e^{\gamma_j m - \frac{\gamma_j^2}{2}} \bar{q}_s(r, m), \quad \bar{\phi}_s^j(r, m) = \frac{1}{2} \int_0^\infty \bar{q}_s(r, y) e^{-\gamma_j y} \alpha_j(m, y) dy.$$

LEMMA 8.5. *Assume $\tilde{x}_j > 0$. We have, for any integer $J \geq 1$,*

$$\lim_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} \left| E_{l, \theta, \xi}^{j, J, n}(s) - \hat{E}_{l, \theta, \xi}^{j, J, n}(s) \right| ds = 0,$$

where

$$\hat{E}_{l, \theta, \xi}^{j, J, n}(s) = \mathbb{E} \left(\int_0^\infty dm \bar{q}_s^j(0, m) \bar{\phi}_{1-s}^j(0, m) \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + R_s^{j, J, n}\}} \right).$$

Proof. We first consider

$$\tilde{E}_{l, \theta, \xi}^{j, J, n}(s) = \mathbb{E} \left(\int_0^\infty dm \bar{q}_{s - \lambda_j J/n}^j(0, m) \bar{\phi}_{1-s - \lambda_j J/n}^j(0, m) \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + R_s^{j, J, n}\}} \right).$$

Let $Z_{l, \theta, \xi}^{j, J, n}(s) = E_{l, \theta, \xi}^{j, J, n}(s) - \tilde{E}_{l, \theta, \xi}^{j, J, n}(s)$ and

$$\zeta_s^j(r_1, r_2, m) = \bar{q}_{s - \lambda_j J/n}^j(r_1, m) \bar{\phi}_{1-s - \lambda_j J/n}^j(r_2, m) - \bar{q}_{s - \lambda_j J/n}^j(0, m) \bar{\phi}_{1-s - \lambda_j J/n}^j(0, m).$$

We have

$$Z_{l, \theta, \xi}^{j, J, n}(s) = \mathbb{E} \left(\int_0^\infty dm \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + R_s^{j, J, n}\}} \zeta_s^j(R_1(\lambda_j J/n), R_2(\lambda_j J/n), m) \right).$$

Now, for all non-negative r_1, r_2, m ,

$$\begin{aligned} |\zeta_s^j(r_1, r_2, m)| &\leq |\bar{q}_{s - \lambda_j J/n}^j(r_1, m) - \bar{q}_{s - \lambda_j J/n}^j(0, m)| |\bar{\phi}_{1-s - \lambda_j J/n}^j(r_2, m)| \\ &\quad + \bar{q}_{s - \lambda_j J/n}^j(0, m) |\bar{\phi}_{1-s - \lambda_j J/n}^j(r_2, m) - \bar{\phi}_{1-s - \lambda_j J/n}^j(0, m)|. \end{aligned}$$

By the arguments of Proposition 6.1, we can easily prove that

$$\left| \frac{\partial \bar{q}_s^j}{\partial r}(r, m) \right| \leq \frac{C}{s^{3/2}} e^{r|\gamma_j|}, \quad \left| \frac{\partial \bar{\phi}_s^j}{\partial r}(r, m) \right| \leq \frac{C}{s} e^{r|\gamma_j| + (\gamma_j^2/2)}.$$

We deduce that

$$|\zeta_s^j(r_1, r_2, m)| \leq C e^{(r_1 + r_2)|\gamma_j| + (\gamma_j^2/2)} r_1 \vee r_2 \delta_j^n(s),$$

with

$$\delta_j^n(s) = \frac{1}{(s - \lambda_j J/n)^{3/2} (1 - s - \lambda_j J/n)^{1/2}} + \frac{1}{(s - \lambda_j J/n) (1 - s - \lambda_j J/n)}.$$

Hence

$$\begin{aligned} \left| Z_{l, \theta, \xi}^{j, J, n}(s) \right| &\leq C \mathbb{E} \left(R_s^{j, J, n} e^{|\gamma_j| (R_1(\lambda_j J/n) + R_2(\lambda_j J/n)) + (\gamma_j^2/2)} \right. \\ &\quad \times R_1(\lambda_j J/n) \vee R_2(\lambda_j J/n) \delta_j^n(s) \Big) \\ &\leq C_{t, \gamma, \sigma} \mathbb{E} \left(R^*(\lambda_j/n) e^{|\gamma_j| (R_1(\lambda_j J/n) + R_2(\lambda_j J/n))} \right. \\ &\quad \times R_1(\lambda_j J/n) \vee R_2(\lambda_j J/n) \delta_j^n(s) \Big) \\ &= C_{t, \gamma, \sigma} \frac{\lambda_j}{n} \mathbb{E} \left(R^*(1) e^{|\gamma_j| \sqrt{\lambda_j/n} (R_1(J) + R_2(J))} R_1(J) \vee R_2(J) \delta_j^n(s) \right), \end{aligned}$$

where the last inequality follows by scaling, and $R^*(u) = R_1^*(u) \vee R_2^*(u)$.

Now, fix $\rho \geq 2$. It can easily be verified that, for n large enough, we have

$$\int_{\frac{\lambda_j \rho J}{n}}^{1 - \frac{\lambda_j \rho J}{n}} \delta_j^n(s) ds \leq C \left(\frac{\sqrt{n}}{\sqrt{\lambda_j(\rho-1)J}} + \log \left(\frac{n}{\lambda_j(\rho-1)J} \right) \right),$$

so that

$$\limsup_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j \rho J}{n}}^{1 - \frac{\lambda_j \rho J}{n}} |Z_{l,\theta,\xi}^{j,J,n}(s)| ds \leq \frac{C_{t,\gamma,\sigma}}{\sqrt{\lambda_j(\rho-1)J}}.$$

On the other hand, it can be proved (as in Lemma 8.2) that, for any fixed $\rho \geq 2$,

$$\int_{\frac{\lambda_j(J+1)}{n}}^{\frac{\lambda_j \rho J}{n}} |Z_{l,\theta,\xi}^{j,J,n}(s)| ds + \int_{1 - \frac{\lambda_j \rho J}{n}}^{1 - \frac{\lambda_j J}{n}} |Z_{l,\theta,\xi}^{j,J,n}(s)| ds = o(1/\sqrt{n}).$$

Therefore

$$\limsup_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} |Z_{l,\theta,\xi}^{j,J,n}(s)| ds \leq \frac{C_{t,\gamma,\sigma}}{\sqrt{\lambda_j(\rho-1)J}},$$

and, by letting ρ go to infinity, we conclude that

$$\lim_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} |E_{l,\theta,\xi}^{j,J,n}(s) - \tilde{E}_{l,\theta,\xi}^{j,J,n}(s)| ds = 0.$$

It remains to show that

$$\lim_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} |\hat{E}_{l,\theta,\xi}^{j,J,n}(s) - \tilde{E}_{l,\theta,\xi}^{j,J,n}(s)| ds = 0.$$

We have, for $s \in \left(\frac{\lambda_j(J+1)}{n}, 1 - \frac{\lambda_j J}{n}\right)$,

$$|\hat{E}_{l,\theta,\xi}^{j,J,n}(s) - \tilde{E}_{l,\theta,\xi}^{j,J,n}(s)| = \left| \mathbb{E} \left(\int_{\tilde{x}_j}^{\tilde{x}_j + R_s^{j,J,n}} dm \eta_j^n(s, m) \right) \right| \leq \mathbb{E} \left(\int_{\tilde{x}_j}^{\tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}} dm |\eta_j^n(s, m)| \right),$$

with

$$\eta_j^n(s, m) = \bar{q}_s^j(0, m) \bar{\phi}_{1-s}^j(0, m) - \bar{q}_{s-\lambda_j J/n}^j(0, m) \bar{\phi}_{1-s-\lambda_j J/n}^j(0, m).$$

Recall that $\bar{q}_s(0, m) = \sqrt{2/\pi} \frac{m}{s^{3/2}} e^{-m^2/2s}$. Note that, if $\tilde{x}_j \leq m \leq \tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}$, we have $e^{-m^2/4s} \leq e^{-\tilde{x}_j^2/4s}$, so that, for $s \in (0, 1)$,

$$\bar{q}_s^j(0, m) \leq \frac{e^{-\tilde{x}_j^2/4s}}{s^{3/2}} m e^{|\gamma_j| m - (m^2/4)} \leq C_j, \quad (8.5)$$

for some positive constant C_j (depending on \tilde{x}_j , but not on s or m).

Furthermore

$$\begin{aligned}\bar{\phi}_s^j(0, m) &= \frac{1}{2} \int_0^\infty dy \bar{q}_s(0, y) e^{-\gamma_j y} \alpha_j(m, y) \\ &= \frac{1}{2\sqrt{s}} \int_0^\infty dy \sqrt{2/\pi} y e^{-y^2/2 - \gamma_j \sqrt{s} y} \alpha_j(m, \sqrt{s} y),\end{aligned}$$

so that $|\bar{\phi}_s^j(0, m)| \leq C/\sqrt{s}$, for some $C > 0$.

We deduce thereof that, for $s \in \left(\frac{\lambda_j(J+1)}{n}, 1 - \frac{\lambda_j J}{n}\right)$,

$$\int_{\tilde{x}_j}^{\tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}} dm |\eta_j^n(s, m)| \leq C_j \frac{R^*(\lambda_j)}{\sqrt{n}} \frac{1}{\sqrt{1 - s - \lambda_j J/n}}.$$

From this estimate, together with the fact that for a fixed s we have $\int_{\tilde{x}_j}^{\tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}} dm |\eta_j^n(s, m)| = o(1/\sqrt{n})$ a.s., we easily deduce that

$$\lim_{n \rightarrow \infty} \sqrt{n} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} |\eta_j^n(s, m)| ds = 0.$$

◇

LEMMA 8.6. *We have, if $\tilde{x}_j > 0$,*

$$\int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} \hat{E}_{l, \theta, \xi}^{j, J, n}(s) ds = \sqrt{\lambda_j/n} \mathbb{E}(R^J) \int_0^1 \varphi_s^j(\tilde{x}_j) ds + o(1/\sqrt{n}),$$

where

$$\varphi_s^j(m) = \bar{q}_s^j(0, m) \bar{\phi}_{1-s}^j(0, m), \quad R^J = \min_{|k| \leq J} \check{R}(U + k),$$

and the random variable U is uniformly distributed on $[0, 1]$ and independent of \check{R} .

Proof. We have

$$\hat{E}_{l, \theta, \xi}^{j, J, n}(s) = \mathbb{E} \int_{\tilde{x}_j}^{\tilde{x}_j + R_s^{j, J, n}} dm \varphi_s^j(m),$$

so that, introducing the notation

$$\bar{E}_{l, \theta, \xi}^{j, J, n}(s) = \mathbb{E}(R_s^{j, J, n}) \bar{q}_s^j(0, \tilde{x}_j) \bar{\phi}_{1-s}^j(0, \tilde{x}_j) = \mathbb{E}(R_s^{j, J, n}) \varphi_s^j(\tilde{x}_j),$$

we can write, for $s \in (\lambda_j(J+1)/n, 1 - \lambda_j J/n)$,

$$\begin{aligned} \left| \hat{E}_{l, \theta, \xi}^{j, J, n}(s) - \bar{E}_{l, \theta, \xi}^{j, J, n}(s) \right| &\leq \mathbb{E} \int_{\tilde{x}_j}^{\tilde{x}_j + R_s^{j, J, n}} dm |\varphi_s^j(m) - \varphi_s^j(\tilde{x}_j)| \\ &\leq \mathbb{E} \int_{\tilde{x}_j}^{\tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}} dm |\varphi_s^j(m) - \varphi_s^j(\tilde{x}_j)|. \end{aligned}$$

Note that, using (8.5) and the estimate $|\bar{\phi}_s^j(0, m)| \leq C/\sqrt{s}$, we have, for $\tilde{x}_j \leq m \leq \tilde{x}_j + R^*(\lambda_j)/\sqrt{n}$,

$$|\varphi_s^j(m)| \leq \frac{C_j}{\sqrt{1-s}}.$$

Now, for a fixed $s \in (0, 1)$, $m \mapsto \varphi_s^j(m)$ is continuous. Indeed the continuity of $\bar{q}_s^j(0, \cdot)$ is obvious and the continuity of $\bar{\phi}_s^j(0, \cdot)$ follows from the continuity assumption on F and the equalities

$$\begin{aligned} \bar{\phi}_s^j(0, m) &= \frac{1}{2} \int_0^\infty dy \bar{q}_s(0, y) e^{-\gamma_j y} \alpha_j(m, y) \\ &= \frac{1}{2} \int_0^\infty dy \bar{q}_s(0, y) e^{-\gamma_j y} F(x_j + \sigma_j(m - y), x_j + \sigma_j m) \\ &= \frac{1}{2} \int_{-\infty}^m dz \bar{q}_s(0, m - z) e^{-\gamma_j(m-z)} F(x_j + \sigma_j z, x_j + \sigma_j m). \end{aligned}$$

Due to the continuity of $m \mapsto \bar{\phi}_s^j(m)$ we have, for a fixed $s \in (0, 1)$,

$$\int_{\tilde{x}_j}^{\tilde{x}_j + \frac{R^*(\lambda_j)}{\sqrt{n}}} dm |\varphi_s^j(m) - \varphi_s^j(\tilde{x}_j)| = o(1/\sqrt{n}) \text{ a.s.,}$$

Hence

$$\int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} \hat{E}_{l, \theta, \xi}^{j, J, n}(s) ds = \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} \mathbb{E}(R_s^{j, J, n}) \varphi_s^j(\tilde{x}_j) ds + o(1/\sqrt{n}).$$

We have

$$\begin{aligned} \mathbb{E}(R_s^{j, J, n}) &= \mathbb{E} \left(\min_{|k| \leq J} \check{R}(d_n^j(s) + \lambda_j k/n) \right) \\ &= \sqrt{\lambda_j/n} \mathbb{E} \left(\min_{|k| \leq J} \check{R}(n d_n^j(s)/\lambda_j + k) \right) \\ &= \sqrt{\lambda_j/n} f \left(\frac{n d_n^j(s)}{\lambda_j} \right), \end{aligned}$$

where, for $u \in [0, 1]$, $f(u) = \mathbb{E} \left(\min_{|k| \leq J} \check{R}(u + k) \right)$. Using the definition of $d_n^j(s)$ and classical arguments, one can show that, for any integrable function g on $[0, 1]$, we have

$$\lim_{n \rightarrow \infty} \int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} f \left(\frac{n d_n^j(s)}{\lambda_j} \right) g(s) ds = \int_0^1 g(s) ds \int_0^1 f(u) du.$$

Therefore

$$\int_{\frac{\lambda_j(J+1)}{n}}^{1 - \frac{\lambda_j J}{n}} \hat{E}_{l, \theta, \xi}^{j, J, n}(s) ds = \sqrt{\lambda_j/n} \mathbb{E}(R^J) \int_0^1 ds \varphi_s^j(\tilde{x}_j) + o(1/\sqrt{n}),$$

which proves the Lemma. \diamond

Proof of Theorem 8.1. Recall that

$$\begin{aligned}\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^{j,n}\}}\right) &= \int_0^1 E_{l,\theta,\xi}^{j,n}(s)ds \\ &= \int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} E_{l,\theta,\xi}^{j,n}(s)ds + o(1/\sqrt{n}),\end{aligned}\quad (8.6)$$

for any positive integer J . Here, we have used Lemma 8.2. Note that, if $x < x_j$, we have, due to Lemma 8.3, $\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^{j,n}\}}\right) = o(1/\sqrt{n})$.

We now assume $x > x_j$ and fix a positive integer J . We have, using Lemma 8.5 and Lemma 8.6,

$$\begin{aligned}\int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} E_{l,\theta,\xi}^{j,J,n}(s)ds &= \int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} \hat{E}_{l,\theta,\xi}^{j,J,n}(s)ds + o(1/\sqrt{n}) \\ &= \sqrt{\frac{\lambda_j}{n}} \mathbb{E}(R^J) \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(1/\sqrt{n}).\end{aligned}$$

Note that $\lim_{J \rightarrow \infty} \mathbb{E}(R^J) = \beta_1$, so that

$$\lim_{J \rightarrow +\infty} \limsup_{n \rightarrow \infty} \sqrt{n} \left(\int_{\frac{\lambda_j(J+1)}{n}}^{1-\frac{\lambda_j J}{n}} E_{l,\theta,\xi}^{j,J,n}(s)ds - \sqrt{\frac{\lambda_j}{n}} \beta_1 \int_0^1 \varphi_s^j(\tilde{x}_j)ds \right) = 0. \quad (8.7)$$

By combining (8.6), (8.7) and Lemma 8.4, we derive

$$\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^{j,n}\}}\right) = \sqrt{\frac{\lambda_j}{n}} \beta_1 \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(1/\sqrt{n}). \quad (8.8)$$

On the other hand, for any $\rho > 0$, we have (using Proposition 5.2 for a function which does not depend on the difference $M^j - M^{j,n}$)

$$\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^j - \rho\}}\right) = \int_0^1 ds \int_0^\infty dm \mathbb{1}_{\{\tilde{x}_j \leq m \leq \tilde{x}_j + \frac{\rho}{\sigma_j}\}} dm \varphi_s^j(m).$$

If $\tilde{x}_j < 0$, we get $\mathbb{E}_{l,\theta,\xi}\left(|F(X_{t_{j+1}^-}, M^j)|\mathbb{1}_{\{M^j \geq x > M^j - \rho\}}\right) = 0$ for $\rho < \sigma_j |\tilde{x}_j|$, so that

$$\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^j - \sigma_{\beta_1} \sqrt{t/n}\}}\right) = o(1/\sqrt{n}).$$

If $\tilde{x}_j > 0$, we have

$$\mathbb{E}_{l,\theta,\xi}\left(|F(X_{t_{j+1}^-}, M^j)|\mathbb{1}_{\{M^j \geq x > M^j - \rho\}}\right) = \frac{\rho}{\sigma_j} \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(\rho)$$

as ρ goes to 0. Therefore

$$\begin{aligned}\mathbb{E}_{l,\theta,\xi}\left(F(X_{t_{j+1}^-}, M^j)\mathbb{1}_{\{M^j \geq x > M^j - \sigma_{\beta_1} \sqrt{t/n}\}}\right) &= \frac{\sigma_{\beta_1} \sqrt{t}}{\sigma_j \sqrt{n}} \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(1/\sqrt{n}) \\ &= \frac{\beta_1 \sqrt{t}}{\sqrt{t_{j+1} - t_j} \sqrt{n}} \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(1/\sqrt{n}) \\ &= \beta_1 \sqrt{\frac{\lambda_j}{n}} \int_0^1 \varphi_s^j(\tilde{x}_j)ds + o(1/\sqrt{n}),\end{aligned}$$

which completes the proof of the second statement of the Theorem. The first one can be proved by the same method. \diamond

REMARK 8.7. It can be deduced from (8.8) that we have an expansion

$$\mathbb{E} \left(g(X_t) \mathbb{1}_{\{M_t \geq x > M_t^n\}} \right) = \frac{C}{\sqrt{n}} + o(1/\sqrt{n}),$$

for some constant C . This can be used to derive an expansion for the difference between continuous and discrete barrier option prices.

REFERENCES

- [1] ASMUSSEN, S., GLYNN P., PITMAN J.: Discretization error in simulation of one-dimensional reflecting brownian motion. *The Annals of Applied Probability* **3**(4), 875-896, (1995).
- [2] BROADIE, M., GLASSERMAN P., KOU S. G.: A Continuity Correction For Discrete Barrier Options. *Mathematical Finance* **7**(4), 325-348, October (1997).
- [3] BROADIE, M., GLASSERMAN P., KOU S. G.: Connecting Discrete and Continuous Path-Dependent Options. *Finance and Stochastics* **3**, 55-82, (1999).
- [4] DIA, E. H. A.: Exotic Options under Exponential Lévy Model. Doctoral thesis, Université Paris-Est, <http://tel.archives-ouvertes.fr/INSMI/tel-00520583/fr/>, (2010).
- [5] DIA, E. H. A., D. LAMBERTON: Connecting Discrete and Continuous Lookback or Hindsight Options in Exponential Lévy Models, *submitted for publication*, 2010.
- [6] KARATZAS, I., SHREVE S.E.: Brownian motion and stochastic calculus, 2nd edition Springer (1991).
- [7] KOU, S. G.: On pricing of discrete barrier options. *Statistica Sinica* **13**, 955-964, (2003).
- [8] KOU, S. G.: A jump-diffusion model for option pricing. *Management Science* **48** (8), 1086-1101, (2002).
- [9] KOU, S. G., WANG H.: Option pricing under a double exponential jump diffusion model. *Management Science* **50** (9), 1178-1192, (2004).
- [10] KYPRIANOU, A. E.: *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer-Verlag, (2006).
- [11] REVUZ, D., YOR M.: *Continuous martingales and Brownian motion*, 2nd edition, Springer, (1994).
- [12] SATO, K.: *Lévy processes and infinitely divisible distributions*. Cambridge University Press, (2005).